

PARTITIONS OF THE WONDERFUL GROUP COMPACTIFICATION

JIANG-HUA LU AND MILEN YAKIMOV

ABSTRACT. We define and study a family of partitions of the wonderful compactification $\overline{G}$ of a semi-simple algebraic group G of adjoint type. The partitions are obtained from subgroups of $G \times G$ associated to triples (A_1, A_2, a) , where A_1 and A_2 are subgraphs of the Dynkin graph Γ of G and $a: A_1 \rightarrow A_2$ is an isomorphism. The partitions of $\overline{G}$ of Springer and Lusztig correspond respectively to the triples $(\emptyset, \emptyset, \text{id})$ and $(\Gamma, \Gamma, \text{id})$.

1. INTRODUCTION

Let G be a connected semi-simple algebraic group over an algebraically closed field k . De Concini and Procesi [4, 5] constructed a wonderful compactification $\overline{G}$ of G , which is a smooth irreducible $(G \times G)$ -variety with finitely many $(G \times G)$ -orbits. Let G_{diag} be the diagonal subgroup of $G \times G$. In his study of parabolic character sheaves on $\overline{G}$ in [14, 15], Lusztig introduced (by an inductive procedure) a partition of $\overline{G}$ by finitely many G_{diag} -stable pieces. The closure of a G_{diag} -stable piece was shown by X. He [7] to be a union of such pieces. Let B be a Borel subgroup of G . Then $\overline{G}$ is also partitioned into finitely many $(B \times B)$ -orbits. The $(B \times B)$ -orbits in $\overline{G}$, as well as their closures, were studied by T. Springer in [19]. In [7], X. He gave a second description of Lusztig's G_{diag} -stable pieces using $(B \times B)$ -orbits in $\overline{G}$, which then enabled him to give [8] an equivalent definition of Lusztig's character sheaves on $\overline{G}$. Further properties and applications of the G_{diag} -stable pieces were obtained by X. He and J. F. Thomsen in [6, 8, 10] and T. A. Springer in [20].

Both G_{diag} and $B \times B$ are special examples of subgroups R_A of $G \times G$ associated to triples (A_1, A_2, a) , where A_1 and A_2 are subgraphs of the Dynkin graph Γ of G , and a is an isomorphism from A_1 to A_2 . If P_{A_1} and P_{A_2} are the standard parabolic subgroups of G corresponding to A_1 and A_2 respectively, then, roughly speaking, R_A is a subgroup of $P_{A_1} \times P_{A_2}$, obtained by identifying the Levi subgroups of P_{A_1} and P_{A_2} via the map a . The precise definition of R_A is given in §2.1. For example, every stabilizer subgroup of $G \times G$ in $\overline{G}$ is conjugate to a group of this form. Moreover, G_{diag} is associated to the triple $(\Gamma, \Gamma, \text{id})$ and $B \times B$ to the triple $(\emptyset, \emptyset, \text{id})$, where $\emptyset$ is the empty set.

In this paper, for any subgroup R_A of $G \times G$ associated to a triple (A_1, A_2, a) , we study a partition of $\overline{G}$ into finitely many R_A -stable pieces indexed by a subset of the Weyl group of $G \times G$. Our definition of the R_A -stable pieces is based on our earlier paper [12] on (R_A, R_C) -double cosets in $G \times G$ for subgroups R_A and R_C associated to any pair of triples (A_1, A_2, a) and (C_1, C_2, c) . We give two additional descriptions of the R_A -stable pieces, which for the case of $R_A = G_{\text{diag}}$, reduce to Lusztig's inductive description in [14, 15] and He's description in [7] using $(B \times B)$ -orbits. In particular, we show that the R_A -stable pieces are smooth, irreducible, locally closed subsets of $\overline{G}$, fibering over flag varieties of Levi subgroups of G . We also show that the closure in $\overline{G}$ of an R_A -stable piece is a union of such pieces. We describe the combinatorics for the closures of the R_A -stable pieces, generalizing both the result of He [7] for the G_{diag} -stable pieces and that of Springer for the $(B \times B)$ -orbits. The closure relations of the R_A -stable pieces are expressed in terms of intersections of the closures with the unique closed $(G \times G)$ -orbit in $\overline{G}$.

Our motivation for studying the $R_{\mathcal{A}}$ -stable pieces in $\overline{G}$ for arbitrary triples (A_1, A_2, a) comes from Poisson geometry. In [13], we study a class of Poisson structures on $\overline{G}$. Special examples of these Poisson structures are induced by the Belavin–Drinfeld r -matrices [1]. The triples (A_1, A_2, a) needed there are precisely the Belavin–Drinfeld triples for the r -matrices. The $R_{\mathcal{A}}$ -stable pieces in $\overline{G}$ as well as their closures are Poisson subvarieties of $\overline{G}$ for the corresponding Poisson structures. To understand these Poisson structures, one needs to first understand the geometry of the $R_{\mathcal{A}}$ -stable pieces.

In §2 – §5, we in fact assume that G_1 and G_2 are any two reductive algebraic groups over an algebraically closed field, and that $R_{\mathcal{A}}$ and $R_{\mathcal{C}}$ are subgroups of $G_1 \times G_2$ associated to two triples (A_1, A_2, a) and (C_1, C_2, c) for $G_1 \times G_2$. The precise definitions of the subgroups $R_{\mathcal{A}}$ and $R_{\mathcal{C}}$ are given in §2.1. Each pair $(R_{\mathcal{A}}, R_{\mathcal{C}})$ of such subgroups gives rise to a decomposition of $G_1 \times G_2$ into $(R_{\mathcal{A}}, R_{\mathcal{C}})$ -stable subsets of the form $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}$, where (v_1, v_2) runs over a subset of the Weyl group for $G_1 \times G_2$ (see (2.4) for detail). In §4, we give a description of the set $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}$ as iterated fiber bundles. The closures of the sets $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}$ in $G_1 \times G_2$ are described in §5. We point out that in a recent paper [9], X. He studies the $(R_{\mathcal{A}}, R_{\mathcal{C}})$ -stable pieces in $G_1 \times G_2$ from the point of view of Coxeter groups. In §6 and §7, the results in §5 are used to prove our main theorems on the $R_{\mathcal{A}}$ -stable pieces in $\overline{G}$ for a semi-simple algebraic group G of adjoint type. Precise statements of our results are summarized in §2. In the appendix we collect a few facts on the Bruhat order on Weyl groups that we use in the paper.

Acknowledgements. We would like to thank Sam Evens and Xuhua He for helpful answers to our questions. We also thank the referees whose comments helped us to improve the exposition.

2. NOTATION AND STATEMENTS OF RESULTS

2.1. Admissible quadruples. For $i = 1, 2$, let G_i be a connected reductive algebraic group over an algebraically closed field k . Let B_i and B_i^- be a fixed pair of opposite Borel subgroups of G_i with U_i being their respective uniradicals. Set $T_i = B_i \cap B_i^-$, and let Γ_i be the set of simple roots determined by (B_i, T_i) . For $\alpha \in \Gamma_i$, denote by U_i^α the one-parameter unipotent subgroup of G_i defined by α . For a subset A_i of Γ_i , let P_{A_i} and $P_{A_i}^-$ be the standard parabolic subgroups of G_i containing respectively B_i and B_i^- . Let $M_{A_i} = P_{A_i} \cap P_{A_i}^-$ be the common Levi factor of $P_{A_i}^\pm$, and let Z_{A_i} be the center of M_{A_i} . The unipotent radicals of P_{A_i} and $P_{A_i}^-$ will be denoted by U_{A_i} and $U_{A_i}^-$ respectively. Let W_i be the Weyl group of Γ_i and W_{A_i} the subgroup of W_i generated by reflections defined by simple roots in A_i . Let $W_i^{A_i}$ and ${}^{A_i}W_i$ be the sets of minimal length representatives of cosets from W_i/W_{A_i} and $W_{A_i} \backslash W_i$ respectively. For each $w_i \in W_i$, we also fix a choice $\tilde{w}_i$ of a representative of w_i in the normalizer of T_i in G_i . The length function on W_i will be denoted by l and the Bruhat order by $\leq$. If a group G acts on a set X , $g.x$ denotes the action of $g \in G$ on $x \in X$. For an element $g \in G$, the map $G \rightarrow G : h \mapsto ghg^{-1}$ will be denoted by Ad_g . The identity element of a group will be denoted by e or 1 .

For subsets A_i of the Dynkin graphs Γ_i , $i = 1, 2$, we call a bijective map $a : A_1 \rightarrow A_2$ an isomorphism, if it preserves the type of each arrow.

Definition 2.1. An admissible quadruple for $G_1 \times G_2$ is a quadruple $\mathcal{A} = (A_1, A_2, a, K)$ consisting of subsets A_1 of Γ_1 and A_2 of Γ_2 , an isomorphism $a : A_1 \rightarrow A_2$, and a closed subgroup K of $M_{A_1} \times M_{A_2}$ of the form

$$(2.1) \quad K = \{(m_1, m_2) \in M_{A_1} \times M_{A_2} \mid \theta_a(m_1 Z_1) = m_2 Z_2\},$$

where, for $i = 1, 2$, Z_i is a closed subgroup of Z_{A_i} and $\theta_a : M_{A_1}/Z_1 \rightarrow M_{A_2}/Z_2$ is an isomorphism mapping T_1/Z_1 to T_2/Z_2 and U_1^α to $U_2^{a(\alpha)}$ for each $\alpha \in A_1$. Here we identify

U_1^α and $U_2^{a(\alpha)}$ with their images in M_{A_1}/Z_1 and M_{A_2}/Z_2 respectively. Given an admissible quadruple $\mathcal{A} = (A_1, A_2, a, K)$ of $G_1 \times G_2$, define

$$(2.2) \quad R_{\mathcal{A}} = K(U_{A_1} \times U_{A_2}) \subset P_{A_1} \times P_{A_2}, \quad R_{\mathcal{A}}^- = K(U_{A_1}^- \times U_{A_2}) \subset P_{A_1}^- \times P_{A_2}.$$

Note that when $G_1 = G_2 = G$, the diagonal subgroup G_{diag} and $B \times B$ for a Borel subgroup B are examples of the groups $R_{\mathcal{A}}$. If further G is of adjoint type, all stabilizer subgroups of $G \times G$ in the De Concini–Procesi [4] compactification $\overline{G}$ of G are conjugate to groups of the type $R_{\mathcal{A}}^-$ (see §2.3).

2.2. An $(R_{\mathcal{A}}, R_{\mathcal{C}})$ -stable partition of $G_1 \times G_2$. In [12] we obtained a classification of $(R_{\mathcal{A}}, R_{\mathcal{C}})$ -double cosets of $G_1 \times G_2$ for two arbitrary admissible quadruples $\mathcal{A} = (A_1, A_2, a, K)$ and $\mathcal{C} = (C_1, C_2, c, L)$ for $G_1 \times G_2$. Given $v_1 \in W_1^{C_1}$, $v_2 \in {}^{A_2}W_2$, set

$$(2.3) \quad C_2(v_1, v_2) = \{\beta \in C_2 \mid (v_2^{-1}av_1c^{-1})^n\beta \text{ is defined and is in } C_2 \text{ for } n = 1, 2, \dots\}.$$

In other words, $C_2(v_1, v_2)$ is the largest subset of C_2 that is stable under $v_2^{-1}av_1c^{-1}$. We proved [12] that each $(R_{\mathcal{A}}, R_{\mathcal{C}})$ -double coset of $G_1 \times G_2$ is of the form $R_{\mathcal{A}}(\dot{v}_1, \dot{v}_2 m_2)R_{\mathcal{C}}$ for some $v_1 \in W_1^{C_1}$, $v_2 \in {}^{A_2}W_2$ and $m_2 \in M_{C_2(v_1, v_2)}$. Moreover, two such double cosets $R_{\mathcal{A}}(\dot{v}_1, \dot{v}_2 m_2)R_{\mathcal{C}}$ and $R_{\mathcal{A}}(\dot{v}'_1, \dot{v}'_2 m'_2)R_{\mathcal{C}}$ coincide if and only if $v'_i = v_i$ for $i = 1, 2$, and m_2 and m'_2 are in the same $(v_2^{-1}av_1c^{-1})$ -twisted conjugacy class in $M_{C_2(v_1, v_2)}$, see Theorem 3.1 for details.

For $v_1 \in W_1^{C_1}$ and $v_2 \in {}^{A_2}W_2$, let

$$(2.4) \quad [v_1, v_2]_{\mathcal{A}, \mathcal{C}} = R_{\mathcal{A}}(v_1, v_2 M_{C_2(v_1, v_2)})R_{\mathcal{C}} \subset G_1 \times G_2.$$

Then by the above result from [12], we have the decomposition

$$(2.5) \quad G_1 \times G_2 = \bigsqcup_{v_1 \in W_1^{C_1}, v_2 \in {}^{A_2}W_2} [v_1, v_2]_{\mathcal{A}, \mathcal{C}}.$$

Here and below $\bigsqcup$ stands for disjoint union. Note that (2.5) is constructed in such a way that the $(R_{\mathcal{A}}, R_{\mathcal{C}})$ -double cosets of $G_1 \times G_2$ corresponding to the same discrete parameters $v_1 \in W_1^{C_1}$ and $v_2 \in {}^{A_2}W_2$ but possibly different continuous parameters $m_2 \in M_{C_2(v_1, v_2)}$ are put together in a single stratum. Alternatively we have the decomposition

$$(2.6) \quad (G_1 \times G_2)/R_{\mathcal{C}} = \bigsqcup_{v_1 \in W_1^{C_1}, v_2 \in {}^{A_2}W_2} [v_1, v_2]_{\mathcal{A}, \mathcal{C}}/R_{\mathcal{C}}$$

of $(G_1 \times G_2)/R_{\mathcal{C}}$ into $R_{\mathcal{A}}$ -stable subsets.

The main objects of study in this paper are the sets $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}$ for $v_1 \in W_1^{C_1}$, $v_2 \in {}^{A_2}W_2$. We describe their geometry, as well as their closure relations. The results are then applied to the wonderful group compactifications.

The following theorem summarizes our results for the decompositions (2.5) and (2.6), see Corollary 4.8, Proposition 4.9, Proposition 4.10, and Theorem 5.2.

Theorem 2.2. *Given any two admissible quadruples $\mathcal{A}$ and $\mathcal{C}$ for $G_1 \times G_2$, the following hold for every $v_1 \in W_1^{C_1}$ and $v_2 \in {}^{A_2}W_2$.*

(i) *$[v_1, v_2]_{\mathcal{A}, \mathcal{C}}$ is locally closed, smooth, and irreducible. Its projection $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}/R_{\mathcal{C}}$ to $(G_1 \times G_2)/R_{\mathcal{C}}$ fibers over the flag variety $M_{A_1}/(M_{A_1} \cap P_{A_1(v_1, v_2)})$ with fibers isomorphic to the product of $M_{C_2(v_1, v_2)}/Y_2$ and the affine space of dimension*

$$\dim U_{A_1(v_1, v_2)} - \dim(U_1 \cap \text{Ad}_{\dot{v}_1} U_{C_1}) + l(v_2),$$

where $A_1(v_1, v_2) = v_1 c^{-1} C_2(v_1, v_2) \subset A_1$, and $Y_2 = \{m \in M_{C_2} \mid (e, m) \in L\} \subset Z_{C_2}$.

(ii) *Alternatively the set $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}$ is given by*

$$(2.7) \quad [v_1, v_2]_{\mathcal{A}, \mathcal{C}} = R_{\mathcal{A}}(B_1 \times B_2)(v_1, v_2)R_{\mathcal{C}}.$$

(iii) The Zariski closure of $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}$ in $G_1 \times G_2$ consists of those $[w_1, w_2]_{\mathcal{A}, \mathcal{C}}$ with $w_1 \in W_1^{C_1}$ and $w_2 \in {}^{A_2}W_2$ for which there exist $x_1 \in W_{A_1}$ and $y_1 \in W_{C_1}$ such that

$$x_1 w_1 y_1 \leq v_1 \quad \text{and} \quad a(x_1) w_2 c(y_1) \leq v_2.$$

(iv) If $\mathcal{A}' = (A_1, A_2, a, K')$ and $\mathcal{C}' = (C_1, C_2, c, L')$ are two other admissible quadruples containing the same triples (A_1, A_2, a) and (C_1, C_2, c) , then there exist $t_2, s_2 \in T_2$ such that

$$[v_1, v_2]_{\mathcal{A}', \mathcal{C}'} = (e, t_2)[v_1, v_2]_{\mathcal{A}, \mathcal{C}}(e, s_2), \quad \forall v_1 \in W_1^{C_1}, v_2 \in {}^{A_2}W_2.$$

Example 2.3. When $\mathcal{A} = \mathcal{C} = (\emptyset, \emptyset, \text{id}, T_1 \times T_2)$ so that $R_{\mathcal{A}} = R_{\mathcal{C}} = B_1 \times B_2$, we have

$$[v_1, v_2]_{\mathcal{A}, \mathcal{C}} = (B_1 \times B_2)(v_1, v_2)(B_1 \times B_2), \quad \forall (v_1, v_2) \in W_1 \times W_2.$$

Thus (2.5) reduces to the Bruhat decomposition

$$G_1 \times G_2 = \bigsqcup_{v_1 \in W_1, v_2 \in W_2} (B_1 \times B_2)(v_1, v_2)(B_1 \times B_2).$$

Part (iii) of Theorem 2.2 in this case is the well-known statement for the closures of Bruhat cells.

2.3. Partitions of the wonderful group compactification. Now we specialize to the case when G_1 and G_2 are both isomorphic to a connected semisimple algebraic group G of adjoint type. All data for G will be denoted as in §2.1, omitting the index i . In particular, B and B^- will be two fixed opposite Borel subgroups of G , $T = B \cap B^-$, and Γ will be the set of simple roots determined by (B, T) .

For $J \subset \Gamma$, let $\pi_J: M_J \rightarrow M_J/Z_J$ be the natural projection. By abuse of notation, we will denote by J the quadruple (J, J, id, L_J) , where

$$L_J = \{(m_1, m_2) \in M_J \times M_J \mid \pi_J(m_1) = \pi_J(m_2)\};$$

so

$$(2.8) \quad R_J^- = L_J(U_J^- \times U_J) = \{(p_1, p_2) \in P_J^- \times P_J \mid \pi_J(p_1) = \pi_J(p_2)\}.$$

Recall that the wonderful compactification $\overline{G}$ of G is a smooth irreducible projective $(G \times G)$ -variety containing G as an open $(G \times G)$ -orbit. It was defined and studied (in the more general framework of symmetric spaces) by De Concini and Procesi [4] in the complex case and by De Concini and Springer [5] for the general case of an algebraically closed field k . The $(G \times G)$ -orbits on $\overline{G}$ are parameterized by the subsets J of Γ . A base point h_J of the $(G \times G)$ -orbit corresponding to $J \subset \Gamma$ has stabilizer subgroup R_J^- .

The partitions (2.5) induce partitions of $\overline{G}$ as follows. Given an arbitrary admissible quadruple $\mathcal{A} = (A_1, A_2, a, K)$ for $G \times G$, by setting

$$(2.9) \quad [J, v_1, v_2]_{\mathcal{A}} = R_{\mathcal{A}}(B \times B)(v_1, v_2).h_J, \quad J \subset \Gamma, v_1 \in W^J, v_2 \in {}^{A_2}W,$$

we obtain the following partition of the wonderful compactification (see §6.1 for detail)

$$(2.10) \quad \overline{G} = \bigsqcup_{J \subset \Gamma, v_1 \in W^J, v_2 \in {}^{A_2}W} [J, v_1, v_2]_{\mathcal{A}},$$

cf. (2.6) and (ii) of Theorem 2.2. We will refer to the sets $[J, v_1, v_2]_{\mathcal{A}}$ in (2.10) as $R_{\mathcal{A}}$ -stable pieces of $\overline{G}$. If the subgroup K in $\mathcal{A}$ is changed to K' , the partition (2.10) is changed by an overall left translation by (e, t) for some $t \in T$ (see (iv) of Theorem 2.2 and Proposition 4.10).

Example 2.4. Let $\mathcal{A}$ be the trivial admissible quadruple $(\emptyset, \emptyset, \text{Id}, T \times T)$. Then $R_{\mathcal{A}} = B \times B$ and we recover Springer's partition [19] of $\overline{G}$ by $(B \times B)$ -orbits:

$$(2.11) \quad \overline{G} = \bigsqcup_{J \subset \Gamma, v_1 \in W^J, v_2 \in W} (B \times B)(v_1, v_2).h_J.$$

On the other hand, let $\mathcal{A}$ be the quadruple $\Gamma_{\text{diag}} := (\Gamma, \Gamma, \text{id}, G_{\text{diag}})$ where G_{diag} is the diagonal subgroup of $G \times G$. By [7], we recover Lusztig's partition [14, 15] of $\overline{G}$:

$$(2.12) \quad \overline{G} = \bigsqcup_{J \subset \Gamma, v_1 \in W^J} [J, v_1, 1]_{\Gamma_{\text{diag}}} = G_{\text{diag}}(B \times B)(v_1, 1).h_J.$$

For a general admissible quadruple $\mathcal{A}$ for $G \times G$, our partition (2.10) is a discrete interpolation between Springer's partition (2.11) and Lusztig's partition (2.12) of $\overline{G}$.

The closure relations of the strata in (2.10) can be derived directly from part (iii) in Theorem 2.2. This is stated in Proposition 7.3. However, we give a more elegant description of the closures using the unique closed $(G \times G)$ -orbit in $\overline{G}$, namely the orbit $(G \times G).h_\emptyset \cong G/B^- \times G/B$, where $\emptyset$ is the empty subset of Γ . More precisely, for any $J \subset \Gamma$ and $v_1 \in W^J$, $v_2 \in {}^{A_2}W$, let $\overline{[J, v_1, v_2]_{\mathcal{A}}}$ be the closure of $[J, v_1, v_2]_{\mathcal{A}}$ in $\overline{G}$, and set

$$\partial_\emptyset \overline{[J, v_1, v_2]_{\mathcal{A}}} = \overline{[J, v_1, v_2]_{\mathcal{A}}} \cap (G \times G).h_\emptyset.$$

The following is a summary of Theorem 7.4 and Theorem 7.6.

Theorem 2.5. *Let $\mathcal{A}$ be an admissible quadruple for $G \times G$. Let $I, J \subset \Gamma$, $v_1 \in W^J$, $v'_1 \in W^I$, and $v_2, v'_2 \in {}^{A_2}W$. Then*

- (i) $[I, v'_1, v'_2]_{\mathcal{A}} \subset \overline{[J, v_1, v_2]_{\mathcal{A}}}$ if and only if $I \subset J$ and $[\emptyset, v'_1, v'_2]_{\mathcal{A}} \subset \partial_\emptyset \overline{[J, v_1, v_2]_{\mathcal{A}}}$;
- (ii) $[I, v'_1, v'_2]_{\mathcal{A}} \subset \overline{[J, v_1, v_2]_{\mathcal{A}}}$ if and only if $I \subset J$ and there exist $x \in W_{A_1}$ and $z \in W_J$ such that $xv'_1 \geq v_1z$ and $a(x)v'_2 \leq v_2z$.

Example 2.6. Consider the case when $\mathcal{A} = (\emptyset, \emptyset, \text{Id}, T \times T)$, so that $R_{\mathcal{A}} = B \times B$. Theorem 2.5 implies that for any $I, J \subset \Gamma$, $v_1 \in W^J$, $v'_1 \in W^I$, and $v_2, v'_2 \in W$,

$$(B \times B)(v'_1, v'_2).h_I \subset \overline{(B \times B)(v_1, v_2).h_J}$$

if and only if $I \subset J$ and there exists $z \in W_J$ such that $v'_1 \geq v_1z$ and $v'_2 \leq v_2z$. This description of the $(B \times B)$ -orbit closures in $\overline{G}$, which is simpler than what is given in [19], was independently obtained in [11]. When $\mathcal{A} = \Gamma_{\text{diag}}$ as in Example 2.4, Theorem 2.5 implies that for any $I, J \subset \Gamma$, $v_1 \in W^J$, $v'_1 \in W^I$,

$$G_{\text{diag}}(B \times B)(v'_1, 1).h_I \subset \overline{G_{\text{diag}}(B \times B)(v_1, 1).h_J}$$

if and only if $I \subset J$ and there exists $x \leq z \in W_J$ such that $xv'_1 \geq v_1z$. It follows from [7, Corollary 3.4] (see Lemma 8.7 in the Appendix) that the above description of the closures of the G_{diag} -stable subsets is the same as that given in [7] by X. He.

3. $(R_{\mathcal{A}}, R_{\mathcal{C}})$ -DOUBLE COSETS IN $G_1 \times G_2$ AND THEIR STABILIZER SUBGROUPS

In this section, we first recall some results from [12].

3.1. Classification of $(R_{\mathcal{A}}, R_{\mathcal{C}})$ -double cosets in $G_1 \times G_2$. Fix two arbitrary admissible quadruples $\mathcal{A} = (A_1, A_2, a, K)$ and $\mathcal{C} = (C_1, C_2, c, L)$ for $G_1 \times G_2$. In [12] we obtained a classification of the $(R_{\mathcal{A}}, R_{\mathcal{C}})$ -double cosets of $G_1 \times G_2$.

For $v_1 \in W_1^{C_1}$ and $v_2 \in {}^{A_2}W_2$, recall the definition (2.3) of the set $C_2(v_1, v_2) \subset C_2$. Similarly, let

$$(3.1) \quad A_1(v_1, v_2) = \{\alpha \in A_1 \mid (v_1 c^{-1} v_2^{-1} a)^n \alpha \text{ is defined and is in } A_1 \text{ for } n = 1, 2, \dots\},$$

so $A_1(v_1, v_2)$ is the largest subset of A_1 that is stable under $v_1 c^{-1} v_2^{-1} a$. Note that

$$(3.2) \quad v_2^{-1} a \text{ and } cv_1^{-1}: A_1(v_1, v_2) \longrightarrow C_2(v_1, v_2)$$

are isomorphisms. Define

$$(3.3) \quad K_{(v_1, v_2)} = (M_{A_1(v_1, v_2)} \times M_{C_2(v_1, v_2)}) \cap \text{Ad}_{(e, \dot{v}_2)}^{-1} K,$$

$$(3.4) \quad L_{(v_1, v_2)} = (M_{A_1(v_1, v_2)} \times M_{C_2(v_1, v_2)}) \cap \text{Ad}_{(\dot{v}_1, e)} L,$$

$$(3.5) \quad Q_{(v_1, v_2)} = \{(m, m') \in M_{C_2(v_1, v_2)} \times M_{C_2(v_1, v_2)} \mid \exists n \in M_{A_1(v_1, v_2)} \text{ such that } (n, m) \in K_{(v_1, v_2)} \text{ and } (n, m') \in L_{(v_1, v_2)}\},$$

and let $Q_{(v_1, v_2)}$ act on $M_{C_2(v_1, v_2)}$ from the left by

$$(3.6) \quad (m, m') \cdot m_2 := mm_2(m')^{-1}, \quad (m, m') \in Q_{(v_1, v_2)}, \quad m_2 \in M_{C_2(v_1, v_2)}.$$

Theorem 3.1. [12] *Let $\mathcal{A} = (A_1, A_2, a, K)$ and $\mathcal{C} = (C_1, C_2, c, L)$ be two admissible quadruples for $G_1 \times G_2$. Then every $(R_{\mathcal{A}}, R_{\mathcal{C}})$ -double coset of $G_1 \times G_2$ is of the form*

$$R_{\mathcal{A}}(\dot{v}_1, \dot{v}_2 m_2) R_{\mathcal{C}} \quad \text{for some } v_1 \in W_1^{C_1}, v_2 \in {}^{A_2}W_2, m_2 \in M_{C_2(v_1, v_2)}.$$

Two such double cosets $R_{\mathcal{A}}(\dot{v}_1, \dot{v}_2 m_2) R_{\mathcal{C}}$ and $R_{\mathcal{A}}(\dot{v}'_1, \dot{v}'_2 m'_2) R_{\mathcal{C}}$ coincide if and only if $v'_i = v_i$ for $i = 1, 2$ and m_2 and m'_2 are in the same $Q_{(v_1, v_2)}$ -orbit in $M_{C_2(v_1, v_2)}$, cf. (3.6).

In [12] we dealt with a slightly more general class of the groups $R_{\mathcal{A}}$ for which the projections $K \rightarrow M_{A_i}$, for $i = 1, 2$, do not have to be surjective.

Recall from (2.4) that for any two admissible quadruples $\mathcal{A}$ and $\mathcal{C}$ for $G_1 \times G_2$,

$$[v_1, v_2]_{\mathcal{A}, \mathcal{C}} = R_{\mathcal{A}}(v_1, v_2 M_{C_2(v_1, v_2)}) R_{\mathcal{C}} \subset G_1 \times G_2$$

for $v_1 \in W_1^{C_1}$ and $v_2 \in {}^{A_2}W_2$. Theorem 3.1 immediately implies:

Corollary 3.2. *For any two admissible quadruples $\mathcal{A}$ and $\mathcal{C}$ for $G_1 \times G_2$,*

$$G_1 \times G_2 = \bigsqcup_{(v_1, v_2) \in W_1^{C_1} \times {}^{A_2}W_2} [v_1, v_2]_{\mathcal{A}, \mathcal{C}} \quad (\text{disjoint union}).$$

Lemma 3.3. *For any $v_1 \in W_1^{C_1}$ and $v_2 \in {}^{A_2}W_2$,*

$$(3.7) \quad [v_1, v_2]_{\mathcal{A}, \mathcal{C}} = R_{\mathcal{A}}(v_1, v_2 (B_2 \cap M_{C_2(v_1, v_2)})) R_{\mathcal{C}},$$

and $R_{\mathcal{A}}(v_1, v_2 T_2) R_{\mathcal{C}}$ is dense in $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}$, where $T_2 = B_2 \cap B_2^-$.

Proof. Let $G_{C_2(v_1, v_2)} = M_{C_2(v_1, v_2)} / Z_{C_2(v_1, v_2)}$. Then

$$\sigma_{(v_1, v_2)} := \text{Ad}_{\dot{v}_2}^{-1} \theta_a \text{Ad}_{\dot{v}_1} \theta_c^{-1}$$

defines an automorphism of $G_{C_2(v_1, v_2)}$. The action of $Q_{(v_1, v_2)}$ on $M_{C_2(v_1, v_2)}$ descends to an action on $G_{C_2(v_1, v_2)}$. The orbits of the latter are exactly the $\sigma_{(v_1, v_2)}$ -twisted conjugacy classes of $G_{C_2(v_1, v_2)}$. (Here and below for a group F and $\sigma \in \text{Aut}(F)$, by a σ -twisted conjugacy class of F we mean an orbit of the action $g.x = gx(\sigma(g))^{-1}$ of F on itself.) Lemma 3.3 now follows from the following results in [16] and [22, Lemma 7.3]: If G is a connected reductive algebraic group, $\sigma \in \text{Aut}(G)$, and B is a σ -stable Borel subgroup G , then

(1) all σ -twisted conjugacy classes of G meet B ;

(2) for every maximal torus T of G inside B the union of all σ -twisted conjugacy classes that meet T is a Zariski open subset of G . $\square$

3.2. Stabilizer subgroups. For $(g_1, g_2) \in G_1 \times G_2$, set

$$(3.8) \quad \text{Stab}_{\mathcal{A}, \mathcal{C}}(g_1, g_2) = R_{\mathcal{A}} \cap \text{Ad}_{(g_1, g_2)} R_{\mathcal{C}}.$$

An explicit description of the subgroups $\text{Stab}_{\mathcal{A}, \mathcal{C}}(g_1, g_2)$ was given in [12]. We only recall some facts about these groups that will be used in this paper (see the proof of Proposition 4.7).

For $v_1 \in W_1^{C_1}$ and $v_2 \in {}^{A_2}W_2$, let $A_2(v_1, v_2) = aA_1(v_1, v_2) \subset A_2$. The following proposition was proved in [12].

Proposition 3.4. *For $q = (\dot{v}_1, \dot{v}_2 m)$, where $v_1 \in W_1^{C_1}$, $v_2 \in {}^{A_2}W_2$, and $m \in M_{C_2(v_1, v_2)}$, $\text{Stab}_{\mathcal{A}, \mathcal{C}}(q) \subset P_{A_1(v_1, v_2)} \times P_{A_2(v_1, v_2)}$, and $\text{Stab}_{\mathcal{A}, \mathcal{C}}(q) = \text{Stab}_{\mathcal{A}, \mathcal{C}}^{\text{red}}(q) \text{Stab}_{\mathcal{A}, \mathcal{C}}^{\text{uni}}(q)$, where*

$$\begin{aligned} \text{Stab}_{\mathcal{A}, \mathcal{C}}^{\text{red}}(q) &= \text{Stab}_{\mathcal{A}, \mathcal{C}}(q) \cap (M_{A_1(v_1, v_2)} \times M_{A_2(v_1, v_2)}) \\ &= (M_{A_1(v_1, v_2)} \times M_{A_2(v_1, v_2)}) \cap K \cap \text{Ad}_{(\dot{v}_1, \dot{v}_2 m)} L \end{aligned}$$

and $\text{Stab}_{\mathcal{A}, \mathcal{C}}^{\text{uni}}(q) = \text{Stab}_{\mathcal{A}, \mathcal{C}}(q) \cap (U_{A_1(v_1, v_2)} \times U_{A_2(v_1, v_2)})$ has dimension equal to

$$\dim(U_1 \cap \text{Ad}_{\dot{v}_1} U_{C_1}) + \dim(U_{A_2} \cap \text{Ad}_{\dot{v}_2} U_2).$$

4. AN INDUCTIVE DESCRIPTION AND THE GEOMETRY OF THE SETS $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}$

In this section, we modify the inductive arguments in [12] to give an inductive description of the sets $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}$. Consequently, we will be able to describe the sets $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}$ as iterated bundles. Our arguments generalize those of Lusztig in [14, 15] for the special case when $G_1 = G_2 = G$ and $R_{\mathcal{A}}$ is the diagonal subgroup of $G \times G$.

4.1. Construction of new admissible quadruples. For $i = 1, 2$ and $E_i, F_i \subset \Gamma_i$, let

$${}^{E_i}W_i^{F_i} = {}^{E_i}W_i \cap W_i^{F_i}.$$

Given an admissible quadruple $\mathcal{C} = (C_1, C_2, c, L)$ for $G_1 \times G_2$, a subset E_1 of Γ_1 , and any $y_1 \in {}^{E_1}W_1^{C_1}$, we construct another admissible quadruple $\mathcal{C}^{(E_1, y_1)}$ for $G_1 \times G_2$ as follows: set

$$\begin{aligned} C_1^{(E_1, y_1)} &= E_1 \cap y_1(C_1), \quad C_2^{(E_1, y_1)} = c(C_1 \cap y_1^{-1}(E_1)) \\ L^{(E_1, y_1)} &= \left(M_{C_1^{(E_1, y_1)}} \times M_{C_2^{(E_1, y_1)}} \right) \cap \text{Ad}_{(\dot{y}_1, e)} L. \end{aligned}$$

Lemma 4.1. *In the above setting the following hold:*

- (i) *The quadruple $\mathcal{C}^{(E_1, y_1)} := (C_1^{(E_1, y_1)}, C_2^{(E_1, y_1)}, c y_1^{-1}, L^{(E_1, y_1)})$ is an admissible quadruple for $G_1 \times G_2$.*
- (ii) *Let $R_{\mathcal{C}^{(E_1, y_1)}}$ be the subgroup of $G_1 \times G_2$ defined by $\mathcal{C}^{(E_1, y_1)}$ as in Definition 2.1. Then $(P_{E_1} \times G_2) \cap \text{Ad}_{(\dot{y}_1, e)} R_{\mathcal{C}} \subset R_{\mathcal{C}^{(E_1, y_1)}}$.*

Proof. Part (i) follows directly from the definition of admissible quadruples. To prove (ii), let $(p, g_2) \in P_{E_1} \times G_2$ be such that $(\dot{y}_1^{-1} p \dot{y}_1, g_2) \in R_{\mathcal{C}}$. For notational simplicity, set

$$D_1 = C_1 \cap y_1^{-1}(E_1) = y_1^{-1}(C_1^{(E_1, y_1)}), \quad D_2 = c(D_1) = C_2^{(E_1, y_1)}.$$

Then by [12, (4.11) in Lemma 4.2],

$$(4.1) \quad \dot{y}_1^{-1} p \dot{y}_1 \in P_{C_1} \cap \text{Ad}_{\dot{y}_1}^{-1} P_{E_1} = M_{D_1}(U_{D_1} \cap \text{Ad}_{\dot{y}_1}^{-1} U_{C_1^{(E_1, y_1)}}) \subset P_{D_1}.$$

It follows from [12, (3) in Lemma 3.4] that $g_2 \in P_{c(D_1)} = P_{D_2} = P_{C_2^{(E_1, y_1)}}$, and

$$\begin{aligned} (\dot{y}_1^{-1} p \dot{y}_1, g_2) &\in (P_{D_1} \times P_{D_2}) \cap R_{\mathcal{C}} = ((P_{D_1} \times P_{D_2}) \cap L)(U_{C_1} \times U_{C_2}) \\ &\subset ((M_{D_1} \times M_{D_2}) \cap L)(U_{D_1} \times U_{D_2}). \end{aligned}$$

Thus by (4.1), $(\dot{y}_1^{-1} p \dot{y}_1, g_2) \in ((M_{D_1} \times M_{D_2}) \cap L)(\text{Ad}_{\dot{y}_1}^{-1} U_{C_1^{(E_1, y_1)}} \times U_{D_2})$ and hence $(p, g_2) \in R_{\mathcal{C}^{(E_1, y_1)}}$. $\square$

4.2. The first step of the induction. Fix two admissible quadruples $\mathcal{A}$ and $\mathcal{C}$ for $G_1 \times G_2$. Given $v_1 \in W_1^{C_1}$ and $v_2 \in {}^{A_2}W_2$, we will use Lemma 4.1 to construct a sequence of admissible quadruples for $G_1 \times G_2$ which will be used to give an inductive description of $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}/R_{\mathcal{C}} \subset (G_1 \times G_2)/R_{\mathcal{C}}$. In this subsection, we present the first step of the inductive description.

Consider the projection

$$\rho_0 : (G_1 \times G_2)/R_{\mathcal{C}} \longrightarrow (G_1 \times G_2)/(P_{C_1} \times P_{C_2}).$$

By [12, Proposition 8.1 and Proposition 4.1], every $R_{\mathcal{A}}$ -orbit in $(G_1 \times G_2)/(P_{C_1} \times P_{C_2})$ contains exactly one point of the form $(x_1, x_2).(P_{C_1} \times P_{C_2})$, where

$$(4.2) \quad x_2 \in {}^{A_2}W_2^{C_2} \quad \text{and} \quad x_1 \in {}^{a^{-1}(A_2 \cap x_2(C_2))}W_1^{C_1}.$$

Let Ω_0 be the set that consists of all pairs (x_1, x_2) satisfying (4.2). For $(x_1, x_2) \in \Omega_0$, let

$$\begin{aligned} \mathcal{O}^{(x_1, x_2)} &= R_{\mathcal{A}}(x_1, x_2).(P_{C_1} \times P_{C_2}) \subset (G_1 \times G_2)/(P_{C_1} \times P_{C_2}), \\ \mathcal{X}^{(x_1, x_2)} &= \rho_0^{-1}(\mathcal{O}^{(x_1, x_2)}) \subset (G_1 \times G_2)/R_{\mathcal{C}}. \end{aligned}$$

Then clearly,

$$\mathcal{X}^{(x_1, x_2)} = \{(r_1 \dot{x}_1, r_2 \dot{x}_2 m).R_{\mathcal{C}} \mid (r_1, r_2) \in R_{\mathcal{A}}, m \in M_{C_2}\}.$$

Letting $E_1 = a^{-1}(A_2 \cap x_2(C_2))$ and $y_1 = x_1$ in Lemma 4.1, we get the admissible quadruple

$$(4.3) \quad \mathcal{C}^{(x_1, x_2)} := \mathcal{C}^{(E_1, x_1)} = (C_1^{(x_1, x_2)}, C_2^{(x_1, x_2)}, c^{(x_1, x_2)}, L^{(x_1, x_2)})$$

and the corresponding subgroup $R_{\mathcal{C}^{(x_1, x_2)}}$ of $G_1 \times G_2$, where in particular,

$$(4.4) \quad C_1^{(x_1, x_2)} = a^{-1}(A_2 \cap x_2(C_2)) \cap x_1(C_1),$$

$$(4.5) \quad C_2^{(x_1, x_2)} = c(C_1 \cap x_1^{-1}a^{-1}(A_2 \cap x_2(C_2))),$$

$$(4.6) \quad c^{(x_1, x_2)} = cx_1^{-1} : C_1^{(x_1, x_2)} \longrightarrow C_2^{(x_1, x_2)}.$$

Lemma 4.2. *For any $(x_1, x_2) \in \Omega_0$, the map $f_0 : \mathcal{X}^{(x_1, x_2)} \rightarrow (G_1 \times G_2)/R_{\mathcal{C}^{(x_1, x_2)}}$ given by*

$$(4.7) \quad f_0 : (r_1 \dot{x}_1, r_2 \dot{x}_2 m).R_{\mathcal{C}} \longmapsto (r_1, r_2 \dot{x}_2 m).R_{\mathcal{C}^{(x_1, x_2)}}, \quad (r_1, r_2) \in R_{\mathcal{A}}, m \in M_{C_2}$$

is a well-defined $R_{\mathcal{A}}$ -equivariant morphism of algebraic varieties.

Proof. Assume that $(r_1 \dot{x}_1, r_2 \dot{x}_2 m).R_{\mathcal{C}} = (r'_1 \dot{x}_1, r'_2 \dot{x}_2 m').R_{\mathcal{C}}$ for $(r_1, r_2), (r'_1, r'_2) \in R_{\mathcal{A}}$ and $m, m' \in M_{C_2}$. Then

$$(4.8) \quad ((r'_1)^{-1}r_1, (r'_2)^{-1}r_2) \in R_{\mathcal{A}} \quad \text{and} \quad (\dot{x}_1^{-1}(r'_1)^{-1}r_1 \dot{x}_1, (m')^{-1}\dot{x}_2^{-1}(r'_2)^{-1}r_2 \dot{x}_2 m) \in R_{\mathcal{C}}.$$

It follows from (4.8) that $(r'_2)^{-1}r_2 \in P_{A_2} \cap x_2(P_{C_2}) \subset P_{A_2 \cap x_2(C_2)}$, which, together with (4.8) and [12, (3) of Lemma 3.4], imply that $(r'_1)^{-1}r_1 \in P_{a^{-1}(A_2 \cap x_2(C_2))}$. By (ii) of Lemma 4.1, $((r'_1)^{-1}r_1, (m')^{-1}\dot{x}_2^{-1}(r'_2)^{-1}r_2 \dot{x}_2 m) \in R_{\mathcal{C}^{(x_1, x_2)}}$. Thus f_0 is well-defined.

Clearly f_0 is $R_{\mathcal{A}}$ -equivariant. To see that f_0 is a morphism, note that since the fiber of ρ_0 over the point $(\dot{x}_1, \dot{x}_2).(P_{C_1} \times P_{C_2})$ is $(\dot{x}_1, \dot{x}_2).(P_{C_1} \times P_{C_2})/R_{\mathcal{C}}$,

$$\mathcal{X}^{(x_1, x_2)} \cong R_{\mathcal{A}} \times_{R_1} (\dot{x}_1, \dot{x}_2).(P_{C_1} \times P_{C_2})/R_{\mathcal{C}},$$

where $R_1 = R_{\mathcal{A}} \cap \text{Ad}_{(\dot{x}_1, \dot{x}_2)}(P_{C_1} \times P_{C_2})$ is the stabilizer subgroup of $R_{\mathcal{A}}$ at the point $(\dot{x}_1, \dot{x}_2).(P_{C_1} \times P_{C_2})$. Let $Y_2 = \{m \in M_{C_2} \mid (e, m) \in L\} \subset Z_{C_2}$. Then the inclusion map of $\{e\} \times M_{C_2} \hookrightarrow P_{C_1} \times P_{C_2}$ induces an isomorphism

$$\phi_1 : (\dot{x}_1, \dot{x}_2)(\{e\} \times M_{C_2})/(\{e\} \times Y_2) \xrightarrow{\cong} (\dot{x}_1, \dot{x}_2).(P_{C_1} \times P_{C_2})/R_{\mathcal{C}}.$$

Note that $\{e\} \times Y_2 \subset L^{(x_1, x_2)} \subset R_{\mathcal{C}^{(x_1, x_2)}}$ so we have the projection

$$\phi_2 : (G_1 \times G_2)/(\{e\} \times Y_2) \rightarrow (G_1 \times G_2)/R_{\mathcal{C}^{(x_1, x_2)}}.$$

Consider the morphism

$$\begin{aligned} \phi_3 : R_{\mathcal{A}} \times (G_1 \times G_2)/(\{e\} \times Y_2) &\longrightarrow (G_1 \times G_2)/(\{e\} \times Y_2), \\ ((r_1, r_2), (g_1, g_2)(\{e\} \times Y_2)) &\longmapsto (r_1 \dot{x}_1^{-1} g_1, r_2 g_2)(\{e\} \times Y_2). \end{aligned}$$

The composition of ϕ_3 with ϕ_2 gives rise to a morphism

$$R_{\mathcal{A}} \times (\dot{x}_1, \dot{x}_2)(\{e\} \times M_{C_2})/(\{e\} \times Y_2) \longrightarrow (G_1 \times G_2)/R_{\mathcal{C}(x_1, x_2)}.$$

Let R_1 act on $(\dot{x}_1, \dot{x}_2)(\{e\} \times M_{C_2})/(\{e\} \times Y_2)$ so that the isomorphism ϕ_1 is R_1 -equivariant. Then the well-definedness of f_0 implies that

$$f_0 : \chi^{(x_1, x_2)} \cong R_{\mathcal{A}} \times_{R_1} (\dot{x}_1, \dot{x}_2)(\{e\} \times M_{C_2})/(\{e\} \times Y_2) \longrightarrow (G_1 \times G_2)/R_{\mathcal{C}(x_1, x_2)}$$

is a morphism of varieties. $\square$

Let now $v_1 \in W_1^{C_1}$ and $v_2 \in {}^{A_2}W_2$. By [3, Proposition 2.7.5] (see Lemma 8.2 in the Appendix), v_2 can be uniquely written as

$$(4.9) \quad v_2 = x_2 u_2, \quad \text{where} \quad x_2 \in {}^{A_2}W_2^{C_2}, \quad \text{and} \quad u_2 \in {}^{C_2 \cap x_2^{-1}(A_2)}W_{C_2}.$$

Given the decomposition $v_2 = x_2 u_2$, write v_1 as

$$(4.10) \quad v_1 = u_1 x_1, \quad \text{where} \quad x_1 \in {}^{a^{-1}(A_2 \cap x_2(C_2))}W_1^{C_1}, \quad \text{and} \quad u_1 \in W_{a^{-1}(A_2 \cap x_2(C_2))}^{C_1^{(x_1, x_2)}}$$

with $C_1^{(x_1, x_2)}$ given in (4.4). Here and below, for $i = 1, 2$ and $E_i \subset D_i \subset \Gamma_i$, we let

$$W_{D_i}^{E_i} = W_{D_i} \cap W_i^{E_i}.$$

It is easy to see that $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}/R_{\mathcal{C}} \subset \chi^{(x_1, x_2)}$. Since $u_1 \in W_{a^{-1}(A_2 \cap x_2(C_2))}^{C_1^{(x_1, x_2)}} \subset W_1^{C_1^{(x_1, x_2)}}$, we have the set

$$[u_1, v_2]_{\mathcal{A}, \mathcal{C}(x_1, x_2)}/R_{\mathcal{C}(x_1, x_2)} = R_{\mathcal{A}} \left(u_1, v_2 M_{C_2^{(x_1, x_2)}(u_1, v_2)} \right) \cdot R_{\mathcal{C}(x_1, x_2)} \subset (G_1 \times G_2)/R_{\mathcal{C}(x_1, x_2)},$$

where $C_2^{(x_1, x_2)}(u_1, v_2)$ is the largest subset of $C_2^{(x_1, x_2)}$ that is stable under the map

$$v_2^{-1} a u_1 (c^{(x_1, x_2)})^{-1} = v_2^{-1} a v_1 c^{-1}.$$

An argument similar to that in the proof of [12, Lemma 5.3] shows that $C_2^{(x_1, x_2)}(u_1, v_2) = C_2(v_1, v_2)$. The following Proposition 4.3 now follows directly from Theorem 3.1.

Proposition 4.3. *Let $v_1 \in W_1^{C_1}$ and $v_2 \in {}^{A_2}W_2$ have the decompositions (4.10) and (4.9). Then $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}$ consists of those $(g_1, g_2) \in G_1 \times G_2$ for which*

$$(g_1, g_2) \cdot R_{\mathcal{C}} \in \mathcal{X}^{(x_1, x_2)} \quad \text{and} \quad f_0((g_1, g_2) \cdot R_{\mathcal{C}}) \in [u_1, v_2]_{\mathcal{A}, \mathcal{C}(x_1, x_2)}/R_{\mathcal{C}(x_1, x_2)}.$$

Moreover, $f_0 : [v_1, v_2]_{\mathcal{A}, \mathcal{C}}/R_{\mathcal{C}} \rightarrow [u_1, v_2]_{\mathcal{A}, \mathcal{C}(x_1, x_2)}/R_{\mathcal{C}(x_1, x_2)}$ induces a bijection between the $R_{\mathcal{A}}$ -orbits in $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}/R_{\mathcal{C}}$ and the $R_{\mathcal{A}}$ -orbits in $[u_1, v_2]_{\mathcal{A}, \mathcal{C}(x_1, x_2)}/R_{\mathcal{C}(x_1, x_2)}$.

4.3. Inductive description of the sets $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}/R_{\mathcal{C}}$. Given $v_1 \in W_1^{C_1}$ and $v_2 \in {}^{A_2}W_2$, by repeating the construction in §4.2, we get a sequence

$$\mathcal{C}^{(i)} = (C_1^{(i)}, C_2^{(i)}, c^{(i)}, L^{(i)}), \quad i \geq 0,$$

of admissible quadruples for $G_1 \times G_2$ and a sequence $u_1^{(i)} \in W_1^{C_1^{(i)}}$, $i \geq 0, \dots$ which gives rise to the sequence of double cosets

$$\mathcal{Z}^{(i)} = [u_1^{(i)}, v_2]_{\mathcal{A}, \mathcal{C}^{(i)}}/R_{\mathcal{C}^{(i)}} \subset (G_1 \times G_2)/R_{\mathcal{C}^{(i)}}, \quad i \geq 0.$$

Here $\mathcal{C}^{(0)} = \mathcal{C}$, $u_1^{(0)} = v_1$, $\mathcal{Z}^{(0)} = [v_1, v_2]_{\mathcal{A}, \mathcal{C}}/R_{\mathcal{C}}$, $\mathcal{C}^{(1)} = \mathcal{C}^{(x_1, x_2)}$ as in (4.3), and $u_1^{(1)} = u_1$ as in (4.10). In general, once $(\mathcal{C}^{(i)}, u_1^{(i)})$ is given, $(\mathcal{C}^{(i+1)}, u_1^{(i+1)})$ is constructed from $(\mathcal{C}^{(i)}, u_1^{(i)})$

in the same way as $(\mathcal{C}^{(x_1, x_2)}, u_1)$ was from $(\mathcal{C}, v_1)$ in §4.2. Namely, first decompose v_2 as the unique product $v_2 = x_2^{(i)} u_2^{(i)}$ with

$$(4.11) \quad x_2^{(i)} \in {}^{A_2}W_2^{C_2^{(i)}}, \quad u_2^{(i)} \in {}^{C_2^{(i)} \cap (x_2^{(i)})^{-1}(A_2)}W_{C_2^{(i)}}.$$

Then decompose $u_1^{(i)}$ as the unique product $u_1^{(i)} = u_1^{(i+1)} x_1^{(i)}$, where

$$x_1^{(i)} \in {}^{a^{-1}(A_2 \cap x_2^{(i)}(C_2^{(i)}))}W_1^{C_1^{(i)}}, \quad u_1^{(i+1)} \in {}^{a^{-1}(A_2 \cap x_2^{(i)}(C_2^{(i)})) \cap x_1^{(i)}(C_1^{(i)})}W_{a^{-1}(A_2 \cap x_2^{(i)}(C_2^{(i)}))}.$$

The admissible quadruple $\mathcal{C}^{(i+1)}$ is constructed as in §4.1 by taking $\mathcal{C}$ to be $\mathcal{C}^{(i)}$, E_1 to be $a^{-1}(A_2 \cap x_2^{(i)}(C_2^{(i)}))$ and y_1 to be $x_1^{(i)}$.

For $i \geq 0$, let ρ_i be the natural projection

$$\rho_i: (G_1 \times G_2)/R_{\mathcal{C}^{(i)}} \longrightarrow (G_1 \times G_2)/(P_{C_1^{(i)}} \times P_{C_2^{(i)}}),$$

$$\mathcal{O}^{(i)} := R_{\mathcal{A}}(x_1^{(i)}, x_2^{(i)}).(P_{C_1^{(i)}} \times P_{C_2^{(i)}}), \text{ and}$$

$$\mathcal{X}^{(i)} := \rho_i^{-1}(\mathcal{O}^{(i)}) \subset (G_1 \times G_2)/R_{\mathcal{C}^{(i)}}.$$

Then $\mathcal{X}^{(i)}$ is a locally closed subset of $(G_1 \times G_2)/R_{\mathcal{C}^{(i)}}$. By Lemma 4.2 and Proposition 4.3, we have a well-defined $R_{\mathcal{A}}$ -equivariant morphism $f_i: \mathcal{X}^{(i)} \longrightarrow (G_1 \times G_2)/R_{\mathcal{C}^{(i+1)}}$ such that

$$(4.12) \quad \mathcal{Z}^{(i)} \subset \mathcal{X}^{(i)} \quad \text{and} \quad \mathcal{Z}^{(i)} = f_i^{-1}(\mathcal{Z}^{(i+1)}), \quad \forall i \geq 0.$$

Let $i_0 \geq 0$ be the smallest integer such that $C_2^{(i_0+1)} = C_2^{(i_0)}$. It is then easy to see that

$$(4.13) \quad \mathcal{C}^{(i_0)} = \mathcal{C}^{(\infty)} := (A_1(v_1, v_2), C_2(v_1, v_2), cv_1^{-1}, L_{(v_1, v_2)}) \quad \text{and} \quad u_1^{(i)} = e, \quad \forall i \geq i_0 + 1,$$

where $L_{(v_1, v_2)}$ is given by (3.4). Set $\mathcal{Z}^{(\infty)} = \mathcal{Z}^{(i_0+1)}$ and $\mathcal{X}^{(\infty)} = \mathcal{X}^{(i_0+1)}$. It follows from (4.13) that

$$\mathcal{Z}^{(i)} = \mathcal{Z}^{(\infty)} = \mathcal{X}^{(\infty)} = [e, v_2]_{\mathcal{A}, \mathcal{C}^{(\infty)}}/R_{\mathcal{C}^{(\infty)}}, \quad \forall i \geq i_0 + 1.$$

Proposition 4.4. *For every $i \geq 0$, $\mathcal{Z}^{(i)}$ is locally closed in $(G_1 \times G_2)/R_{\mathcal{C}^{(i)}}$.*

Proof. By the inductive description of the sets $\mathcal{Z}^{(i)}$ in (4.12), if for some $i \geq 0$, $\mathcal{Z}^{(i+1)}$ is locally closed in $(G_1 \times G_2)/R_{\mathcal{C}^{(i+1)}}$, then $\mathcal{Z}^{(i)}$ is locally closed in $(G_1 \times G_2)/R_{\mathcal{C}^{(i)}}$ because f_i is a morphism. Since $\mathcal{Z}^{(\infty)} = \mathcal{X}^{(\infty)}$ is locally closed in $(G_1 \times G_2)/R_{\mathcal{C}^{(\infty)}}$, it follows that $\mathcal{Z}^{(i)}$ is locally closed in $(G_1 \times G_2)/R_{\mathcal{C}^{(i)}}$ for every $i \geq 0$. $\square$

Let $\rho_{\infty} := \rho_{i_0+1}: (G_1 \times G_2)/R_{\mathcal{C}^{(\infty)}} \rightarrow (G_1 \times G_2)/(P_{A_1(v_1, v_2)} \times P_{C_2(v_1, v_2)})$, and

$$\mathcal{O}^{(\infty)} := \mathcal{O}^{(i_0+1)} = R_{\mathcal{A}}(e, v_2).(P_{A_1(v_1, v_2)} \times P_{C_2(v_1, v_2)}),$$

so that $\mathcal{Z}^{(\infty)} = \mathcal{X}^{(\infty)} = \rho_{\infty}^{-1}(\mathcal{O}^{(\infty)})$. Then we have the projection $\rho_{\infty}: \mathcal{Z}^{(\infty)} \rightarrow \mathcal{O}^{(\infty)}$. Note that the stabilizer subgroup of $R_{\mathcal{A}}$ at the point

$$(e, v_2).(P_{A_1(v_1, v_2)} \times P_{C_2(v_1, v_2)}) \in (G_1 \times G_2)/(P_{A_1(v_1, v_2)} \times P_{C_2(v_1, v_2)})$$

is

$$(4.14) \quad R_{\mathcal{A}} \cap (P_{A_1(v_1, v_2)} \times \text{Ad}_{v_2} P_{C_2(v_1, v_2)}) \subset R_{\mathcal{A}} \cap (P_{A_1(v_1, v_2)} \times P_{A_2(v_1, v_2)}),$$

where $A_2(v_1, v_2) = aA_1(v_1, v_2)$. For notational simplicity, set

$$(4.15) \quad R_{(v_1, v_2)} = R_{\mathcal{A}} \cap (P_{A_1(v_1, v_2)} \times P_{A_2(v_1, v_2)}).$$

Let $\rho'_{\infty}: \mathcal{O}^{(\infty)} \rightarrow R_{\mathcal{A}}/R_{(v_1, v_2)}$ be the projection induced by the inclusion map in (4.14). Then we have the sequence of $R_{\mathcal{A}}$ -equivariant morphisms

$$(4.16) \quad [v_1, v_2]_{\mathcal{A}, \mathcal{C}}/R_{\mathcal{C}} = \mathcal{Z}^{(0)} \xrightarrow{f_0} \mathcal{Z}^{(1)} \xrightarrow{f_1} \dots \xrightarrow{f_{i_0-1}} \mathcal{Z}^{(i_0)} \xrightarrow{f_{i_0}} \mathcal{Z}^{(\infty)} \xrightarrow{\rho_{\infty}} \mathcal{O}^{(\infty)} \xrightarrow{\rho'_{\infty}} R_{\mathcal{A}}/R_{(v_1, v_2)}.$$

Lemma 4.5. *The quotient $R_{\mathcal{A}}/R_{(v_1, v_2)}$ is isomorphic to the flag variety $M_{A_1}/(M_{A_1} \cap P_{A_1(v_1, v_2)})$ of M_{A_1} . The fibers of $\rho_{\infty} : \mathcal{Z}^{(\infty)} \rightarrow \mathcal{O}^{(\infty)}$ are isomorphic to $M_{C_2(v_1, v_2)}/Y_2$, where $Y_2 = \{m \in M_{C_2} \mid (e, m) \in L\} \subset Z_{C_2}$, and the fibers of $\rho'_{\infty} : \mathcal{O}^{(\infty)} \rightarrow R_{\mathcal{A}}/R_{(v_1, v_2)}$ are isomorphic to $U_2 \cap \text{Ad}_{\dot{v}_2} U_2^-$.*

Proof. Consider the group homomorphism

$$p: R_{\mathcal{A}} \longrightarrow M_{A_1}: (r_1, r_2) \longmapsto m_1 \quad \text{if } r_1 = m_1 u_1 \text{ for } m_1 \in M_{A_1}, u_1 \in U_{A_1}.$$

Since p is surjective, the action of $R_{\mathcal{A}}$ on $M_{A_1}/(M_{A_1} \cap P_{A_1(v_1, v_2)})$ through the homomorphism p is transitive. The stabilizer subgroup of $R_{\mathcal{A}}$ at the point $e \cdot (M_{A_1} \cap P_{A_1(v_1, v_2)})$ is $R_{(v_1, v_2)}$ by [12, (3.7) in Lemma 3.4]. Thus $R_{\mathcal{A}}/R_{(v_1, v_2)}$ is isomorphic to $M_{A_1}/(M_{A_1} \cap P_{A_1(v_1, v_2)})$.

The fibers of $\rho_{\infty} : \mathcal{Z}^{(\infty)} \rightarrow \mathcal{O}^{(\infty)}$ are clearly isomorphic to

$$(P_{A_1(v_1, v_2)} \times P_{C_2(v_1, v_2)})/R_{\mathcal{C}^{(\infty)}} \cong (M_{A_1(v_1, v_2)} \times M_{C_2(v_1, v_2)})/L_{(v_1, v_2)}.$$

Note that $\{e\} \times M_{C_2(v_1, v_2)}$ acts on $(M_{A_1(v_1, v_2)} \times M_{C_2(v_1, v_2)})/L_{(v_1, v_2)}$ transitively, and

$$(\{e\} \times M_{C_2(v_1, v_2)}) \cap L_{(v_1, v_2)} \cong \{e\} \times Y_2.$$

Thus the fibers of $\rho_{\infty} : \mathcal{Z}^{(\infty)} \rightarrow \mathcal{O}^{(\infty)}$ are isomorphic to $M_{C_2(v_1, v_2)}/Y_2$. The fibers of ρ'_{∞} are isomorphic to

$$R_{(v_1, v_2)}/R_{\mathcal{A}} \cap (P_{A_1(v_1, v_2)} \times \text{Ad}_{\dot{v}_2} P_{C_2(v_1, v_2)}) \cong U_2 \cap \text{Ad}_{\dot{v}_2} U_2^-.$$

□

4.4. The set $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}/R_{\mathcal{C}}$ as an iterated fiber bundle. Assume the setting from §4.3. We will show in this subsection that each morphism $f_i : \mathcal{Z}^{(i)} \rightarrow \mathcal{Z}^{(i+1)}$ in (4.16) has fibers isomorphic to an affine space.

Fix $m \in M_{C_2(v_1, v_2)}$. For $i \geq 0$, let $z_m^{(i)} = (\dot{u}_1^{(i)}, \dot{v}_2 m) \cdot R_{\mathcal{C}^{(i)}} \in \mathcal{Z}^{(i)}$ and

$$S_m^{(i)} = R_{\mathcal{A}} \cap \text{Ad}_{(\dot{u}_1^{(i)}, \dot{v}_2 m)} R_{\mathcal{C}^{(i)}}$$

be the stabilizer subgroup of $R_{\mathcal{A}}$ at $z_m^{(i)} \in \mathcal{Z}^{(i)}$. Set

$$F_m^{(i)} = f_i^{-1}(z_m^{(i+1)}) \subset \mathcal{Z}^{(i)}.$$

Lemma 4.6. *For any $m \in M_{C_2(v_1, v_2)}$ and $i \geq 0$, $S_m^{(i)} \subset S_m^{(i+1)}$,*

$$(4.17) \quad F_m^{(i)} = S_m^{(i+1)} \cdot z_m^{(i)} \cong S_m^{(i+1)}/S_m^{(i)}.$$

Proof. Let $i \geq 0$. Since $f_i(z_m^{(i)}) = z_m^{(i+1)}$ and since f_i is $R_{\mathcal{A}}$ -equivariant, we have $S_m^{(i)} \subset S_m^{(i+1)}$, and $S_m^{(i+1)} \cdot z_m^{(i)} \subset F_m^{(i)}$. It suffices to prove (4.17) for $i = 0$, and we only need to show that $F_m^{(0)} \subset S_m^{(1)} \cdot z_m^{(0)}$.

Recall that $\mathcal{C}^{(1)} = \mathcal{C}^{(x_1, x_2)}$ as in (4.3), and $u^{(1)} = u_1$ as in the $v_1 = u_1 x_1$ decomposition in (4.10). To show that $F_m^{(0)} \subset S_m^{(1)} \cdot z_m^{(0)}$, assume that $(r_1, r_2) \in R_{\mathcal{A}}$ and $m' \in M_{C_2(v_1, v_2)}$ are such that $(r_1 \dot{v}_1, r_2 \dot{v}_2 m') \cdot R_{\mathcal{C}} \in F_m^{(0)}$. Then

$$R_{\mathcal{A}}(\dot{u}_1, \dot{v}_2 m') \cdot R_{\mathcal{C}^{(x_1, x_2)}} = R_{\mathcal{A}}(\dot{u}_1, \dot{v}_2 m) \cdot R_{\mathcal{C}^{(x_1, x_2)}}.$$

Applying Theorem 3.1 to the quadruples $(\mathcal{A}, \mathcal{C}^{(x_1, x_2)})$ and noting that the largest subset of A_1 that is stable under $u_1(c^{(x_1, x_2)})^{-1} v_2^{-1} a = v_1 c^{-1} v_2^{-1} a$ is $A_1(v_1, v_2)$ and that

$$(M_{A_1(v_1, v_2)} \times M_{C_2(v_1, v_2)}) \cap \text{Ad}_{(\dot{u}_1, e)} L^{(x_1, x_2)} = L_{(v_1, v_2)},$$

we know that there exist $m_1 \in M_{A_1(v_1, v_2)}$ and $n, n' \in M_{C_2(v_1, v_2)}$ with $(m_1, n) \in K_{(v_1, v_2)}$ and $(m_1, n') \in L_{(v_1, v_2)}$ such that $m' = n m (n')^{-1}$, where $K_{(v_1, v_2)}$ and $L_{(v_1, v_2)}$ are respectively

given by (3.3) and (3.4). Thus

$$\begin{aligned} (r_1 \dot{v}_1, r_2 \dot{v}_2 m').R_{\mathcal{C}} &= (r_1 \dot{v}_1, r_2 \dot{v}_2 n m(n')^{-1}).R_{\mathcal{C}} = (r_1 m_1 \dot{v}_1, r_2 \dot{v}_2 n m).R_{\mathcal{C}} \\ &= (r_1, r_2)(m_1, \text{Ad}_{\dot{v}_2}(n))(\dot{v}_1, \dot{v}_2 m).R_{\mathcal{C}} = (r'_1, r'_2)z_m^{(0)}, \end{aligned}$$

where $(r'_1, r'_2) = (r_1, r_2)(m_1, \text{Ad}_{\dot{v}_2}(n)) \in R_{\mathcal{A}}$. On the other hand, it follows from $(r_1 \dot{v}_1, r_2 \dot{v}_2 m').R_{\mathcal{C}} \in F_m^{(0)}$ that

$$z_m^{(1)} = (r_1 \dot{u}_1, r_2 \dot{v}_2 m').R_{\mathcal{C}^{(x_1, x_2)}} = (r_1 \dot{u}_1, r_2 \dot{v}_2 n m(n')^{-1}).R_{\mathcal{C}^{(x_1, x_2)}} = (r'_1, r'_2)z_m^{(1)},$$

so $(r'_1, r'_2) \in S_m^{(1)}$, and thus $(r_1 \dot{v}_1, r_2 \dot{v}_2 m').R_{\mathcal{C}} \in S_m^{(1)}.z_m^{(0)}$. Hence $F_m^{(0)} \subset S_m^{(1)}.z_m^{(0)}$. $\square$

Proposition 4.7. *For each $i \geq 0$, the fibers of the morphism $f_i : \mathcal{Z}^{(i)} \rightarrow \mathcal{Z}^{(i+1)}$ are isomorphic to the affine space of dimension*

$$(4.18) \quad \dim \left(U_1 \cap \text{Ad}_{u_1^{(i+1)}} U_{C_1^{(i+1)}} \right) - \dim \left(U_1 \cap \text{Ad}_{u_1^{(i)}} U_{C_1^{(i)}} \right).$$

Proof. Let $i \geq 0$ and $m \in M_{C_2(v_2, v_2)}$. By Proposition 3.4, we have the semi-direct product decompositions $S_m^{(i)} = S_m^{(i), \text{red}} S_m^{(i), \text{uni}}$, where

$$S_m^{(i), \text{red}} = S_m^{(i)} \cap (M_{A_1(v_1, v_2)} \times M_{A_2(v_1, v_2)}), \quad S_m^{(i), \text{uni}} = S_m^{(i)} \cap (U_{A_1(v_1, v_2)} \times U_{A_2(v_1, v_2)}).$$

It also follows from Proposition 3.4 that $S_m^{(i), \text{red}} = S_m^{(i+1), \text{red}}$. Thus by (4.17) and by Proposition 3.4, $F_m^{(i)} \cong S_m^{(i+1), \text{uni}} / S_m^{(i), \text{uni}}$ is isomorphic to an affine space of dimension (4.18). $\square$

For $v_1 \in W_1^{C_1}$ and $v_2 \in {}^{A_2}W_2$, consider the composition

$$p_0 = \rho'_\infty \circ \rho_\infty \circ f_{i_0} \circ f_{i_0-1} \circ \cdots \circ f_1 \circ f_0 : [v_1, v_2]_{\mathcal{A}, \mathcal{C}} \longrightarrow R_{\mathcal{A}} / R_{(v_1, v_2)}.$$

Since for $(r_1, r_2) \in R_{\mathcal{A}}$ and $m \in M_{C_2(v_1, v_2)}$,

$$p_0((r_1, r_2)(\dot{v}_1, \dot{v}_2 m).R_{\mathcal{C}}) = (r_1, r_2).R_{(v_1, v_2)},$$

it follows that the fiber of p_0 over the point $e.R_{(v_1, v_2)} \in R_{\mathcal{A}} / R_{(v_1, v_2)}$ is

$$X_{(v_1, v_2)} \stackrel{\text{def}}{=} R_{(v_1, v_2)}(v_1, v_2 M_{C_2(v_1, v_2)}).R_{\mathcal{C}} \subset [v_1, v_2]_{\mathcal{A}, \mathcal{C}} / R_{\mathcal{C}}.$$

Introduce

$$P'_{A_2(v_1, v_2)} = M_{A_2(v_1, v_2)}(U_{A_2(v_1, v_2)} \cap \text{Ad}_{\dot{v}_2} U_2^-) \subset P_{A_2(v_1, v_2)}.$$

Since $M_{A_2(v_1, v_2)}$ normalizes $U_{A_2(v_1, v_2)} \cap \text{Ad}_{\dot{v}_2} U_2^-$, $P'_{A_2(v_1, v_2)}$ is a subgroup of $P_{A_2(v_1, v_2)}$.

Corollary 4.8. *For any $v_1 \in W_1^{C_1}$ and $v_2 \in {}^{A_2}W_2$, (i) $X_{(v_1, v_2)}$ is isomorphic to*

$$\begin{aligned} (P_{A_1(v_1, v_2)} \times P_{A_2(v_1, v_2)})(\dot{v}_1, \dot{v}_2).R_{\mathcal{C}} &= (U_{A_1(v_1, v_2)} \times P'_{A_2(v_1, v_2)})(\dot{v}_1, \dot{v}_2).R_{\mathcal{C}} \\ &\cong (U_{A_1(v_1, v_2)} / (U_{A_1(v_1, v_2)} \cap \text{Ad}_{\dot{v}_1} U_{C_1})) \times (U_2 \cap \text{Ad}_{\dot{v}_2} U_2^-) \times (M_{C_2(v_1, v_2)} / Y_2). \end{aligned}$$

In particular, $X_{(v_1, v_2)}$ is a smooth locally closed subset of $(G_1 \times G_2) / R_{\mathcal{C}}$;

(ii) the action map of $R_{\mathcal{A}}$ on $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}$ gives rise to an isomorphism

$$R_{\mathcal{A}} \times_{R_{(v_1, v_2)}} X_{(v_1, v_2)} \longrightarrow [v_1, v_2]_{\mathcal{A}, \mathcal{C}} / R_{\mathcal{C}}.$$

In particular, $[v_1, v_2]_{\mathcal{A}, \mathcal{C}} / R_{\mathcal{C}}$ is a smooth irreducible locally closed subset of $(G_1 \times G_2) / R_{\mathcal{C}}$.

Since $G_1 \times G_2 \rightarrow (G_1 \times G_2) / R_{\mathcal{C}}$ is a locally trivial fibration, $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}$ is a smooth irreducible locally closed subset of $G_1 \times G_2$. Note also that $U_{A_1(v_1, v_2)} \cap \text{Ad}_{\dot{v}_1} U_{C_1} = U_1 \cap \text{Ad}_{\dot{v}_1} U_{C_1}$, so (i) of Theorem 2.2 follows now from Corollary 4.8.

Proof. (i) Clearly $X_{(v_1, v_2)} \subset (P_{A_1(v_1, v_2)} \times P_{A_2(v_1, v_2)})(\dot{v}_1, \dot{v}_2).R_{\mathcal{C}}$. Since

$$\begin{aligned} P_{A_1(v_1, v_2)} \times P_{A_2(v_1, v_2)} &= R_{(v_1, v_2)}(\{e\} \times P_{A_2(v_1, v_2)} \cap M_{A_2}) \\ &= R_{(v_1, v_2)}(\{e\} \times M_{A_2(v_1, v_2)}(U_{A_2(v_1, v_2)} \cap M_{A_2})), \end{aligned}$$

we have

$$(P_{A_1(v_1, v_2)} \times P_{A_2(v_1, v_2)})(\dot{v}_1, \dot{v}_2).R_{\mathcal{C}} = R_{(v_1, v_2)}(\dot{v}_1, \dot{v}_2 M_{C_2(v_1, v_2)} \text{Ad}_{\dot{v}_2}^{-1}(U_{A_2(v_1, v_2)} \cap M_{A_2})).R_{\mathcal{C}}.$$

Since $v_2 \in {}^{A_2}W_2$, $\text{Ad}_{\dot{v}_2}^{-1}(U_{A_2(v_1, v_2)} \cap M_{A_2}) \subset U_2$. Thus

$$(P_{A_1(v_1, v_2)} \times P_{A_2(v_1, v_2)})(\dot{v}_1, \dot{v}_2).R_{\mathcal{C}} \subset R_{(v_1, v_2)}(\dot{v}_1, \dot{v}_2 P_{C_2(v_1, v_2)}).R_{\mathcal{C}}.$$

By following the definition of f_i for each $i \geq 0$, one sees that $(\dot{v}_1, \dot{v}_2 P_{C_2(v_1, v_2)}).R_{\mathcal{C}} \subset X_{(v_1, v_2)}$. Thus $(P_{A_1(v_1, v_2)} \times P_{A_2(v_1, v_2)})(\dot{v}_1, \dot{v}_2).R_{\mathcal{C}} = X_{(v_1, v_2)}$. Now since

$$\begin{aligned} P_{A_2(v_1, v_2)} &= M_{A_2(v_1, v_2)}(U_{A_2(v_1, v_2)} \cap \text{Ad}_{\dot{v}_2} U_2^-)(U_{A_2(v_1, v_2)} \cap \text{Ad}_{\dot{v}_2} U_2) \\ &= P'_{A_2(v_1, v_2)}(U_{A_2(v_1, v_2)} \cap \text{Ad}_{\dot{v}_2} U_2), \end{aligned}$$

we have $P_{A_2(v_1, v_2)} \dot{v}_2 = P'_{A_2(v_1, v_2)} \dot{v}_2 (U_2 \cap \text{Ad}_{\dot{v}_2}^{-1} U_{A_2(v_1, v_2)})$. Thus

$$X_{(v_1, v_2)} = (P_{A_1(v_1, v_2)} \dot{v}_1, P'_{A_2(v_1, v_2)} \dot{v}_2 (U_2 \cap \text{Ad}_{\dot{v}_2}^{-1} U_{A_2(v_1, v_2)} \cap M_{C_2})).R_{\mathcal{C}}.$$

Using the facts that $v_2^{-1} A_2(v_1, v_2) = C_2(v_1, v_2)$ and $v_1 c^{-1} C_2(v_1, v_2) = A_1(v_1, v_2)$, we have

$$X_{(v_1, v_2)} = (P_{A_1(v_1, v_2)} \dot{v}_1, P'_{A_2(v_1, v_2)} \dot{v}_2).R_{\mathcal{C}}.$$

Using again the fact that $c v_1^{-1} A_1(v_1, v_2) = C_2(v_1, v_2)$, we get

$$X_{(v_1, v_2)} = (U_{A_1(v_1, v_2)} \times P'_{A_2(v_1, v_2)})(\dot{v}_1, \dot{v}_2).R_{\mathcal{C}}.$$

Let $S = (U_{A_1(v_1, v_2)} \times P'_{A_2(v_1, v_2)}) \cap \text{Ad}_{(\dot{v}_1, \dot{v}_2)} R_{\mathcal{C}}$. We now determine S . Suppose that $u \in U_{A_1(v_1, v_2)}$ and $p' \in P'_{A_2(v_1, v_2)}$ are such that $(u, p') \in S$. Then $(\dot{v}_1^{-1} u \dot{v}_1, \dot{v}_2^{-1} p' \dot{v}_2) \in R_{\mathcal{C}}$. Hence

$$\dot{v}_1^{-1} u \dot{v}_1 \in (\text{Ad}_{\dot{v}_1}^{-1} U_{A_1(v_1, v_2)}) \cap P_{C_1} = (\text{Ad}_{\dot{v}_1}^{-1} U_{A_1(v_1, v_2)}) \cap U_{C_1}.$$

Thus $\dot{v}_2^{-1} p' \dot{v}_2 \in Y_2 U_{C_2}$. Since $\dot{v}_2^{-1} p' \dot{v}_2 \in M_{C_2(v_1, v_2)} U_2^- \subset P_{C_2(v_1, v_2)}^-$, we must have $\dot{v}_2^{-1} p' \dot{v}_2 \in Y_2$. Thus $(u, p') \in (U_{A_1(v_1, v_2)} \cap \text{Ad}_{\dot{v}_1} U_{C_1}) \times \text{Ad}_{\dot{v}_2} Y_2$. Since

$$(U_{A_1(v_1, v_2)} \cap \text{Ad}_{\dot{v}_1} U_{C_1}) \times \text{Ad}_{\dot{v}_2} Y_2 \subset S,$$

we have $S = (U_{A_1(v_1, v_2)} \cap \text{Ad}_{\dot{v}_1} U_{C_1}) \times \text{Ad}_{\dot{v}_2} Y_2$. Thus $X_{(v_1, v_2)}$ is isomorphic to

$$\begin{aligned} &(U_{A_1(v_1, v_2)} \times P'_{A_2(v_1, v_2)})/S \\ &\cong (U_{A_1(v_1, v_2)} / (U_{A_1(v_1, v_2)} \cap \text{Ad}_{\dot{v}_1} U_{C_1})) \times (U_{A_2(v_1, v_2)} \cap \text{Ad}_{\dot{v}_2} U_2^-) \times (M_{A_2(v_1, v_2)} / \text{Ad}_{\dot{v}_2} Y_2) \\ &\cong (U_{A_1(v_1, v_2)} / (U_{A_1(v_1, v_2)} \cap \text{Ad}_{\dot{v}_1} U_{C_1})) \times (U_2 \cap \text{Ad}_{\dot{v}_2} U_2^-) \times (M_{C_2(v_1, v_2)} / Y_2). \end{aligned}$$

(ii) Since p_0 is $R_{\mathcal{A}}$ -equivariant, by [17, Lemma 4], the action map of $R_{\mathcal{A}}$ on $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}$ induces an isomorphism

$$R_{\mathcal{A}} \times_{R_{(v_1, v_2)}} X(v_1, v_2) \longrightarrow [v_1, v_2]_{\mathcal{A}, \mathcal{C}}$$

of $R_{\mathcal{A}}$ -varieties. $\square$

4.5. Another description of the strata $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}/R_{\mathcal{C}}$.

Proposition 4.9. *For any $v_1 \in W_1^{C_1}$, $v_2 \in {}^{A_2}W_2$, one has*

$$(4.19) \quad [v_1, v_2]_{\mathcal{A}, \mathcal{C}}/R_{\mathcal{C}} = R_{\mathcal{A}}(B_1 \times B_2)(v_1, v_2).R_{\mathcal{C}}.$$

Proof. By Lemma 3.3, we have

$$\begin{aligned} [v_1, v_2]_{\mathcal{A}, \mathcal{C}}/R_{\mathcal{C}} &= R_{\mathcal{A}}(v_1, v_2(B_2 \cap M_{C_2(v_1, v_2)})).R_{\mathcal{C}} \subset \\ &\quad R_{\mathcal{A}}(v_1, v_2 B_2).R_{\mathcal{C}} = R_{\mathcal{A}}(B_1 \times B_2)(v_1, v_2).R_{\mathcal{C}}. \end{aligned}$$

It remains to show that $R_{\mathcal{A}}(B_1 \times B_2)(v_1, v_2).R_{\mathcal{C}} \subset [v_1, v_2]_{\mathcal{A}, \mathcal{C}}/R_{\mathcal{C}}$. Write $v_2 = x_2 u_2$ and $v_1 = u_1 x_1$ as in (4.9) and (4.10) and let $f_0: \mathcal{X}^{(x_1, x_2)} \rightarrow (G_1 \times G_2)/R_{\mathcal{C}(x_1, x_2)}$ be as in Lemma 4.2. Then

$$\begin{aligned} R_{\mathcal{A}}(B_1 \times B_2)(v_1, v_2).R_{\mathcal{C}} &= R_{\mathcal{A}}(v_1, v_2(B_2 \cap M_{C_2})).R_{\mathcal{C}} \subset \mathcal{X}^{(x_1, x_2)}, \\ f_0(R_{\mathcal{A}}(B_1 \times B_2)(v_1, v_2).R_{\mathcal{C}}) &= R_{\mathcal{A}}(u_1, v_2(B_2 \cap M_{C_2})).R_{\mathcal{C}(x_1, x_2)} \\ &= R_{\mathcal{A}}(u_1, v_2(B_2 \cap M_{C_2^{(x_1, x_2)}})).R_{\mathcal{C}(x_1, x_2)} \\ &= R_{\mathcal{A}}(B_1 \times B_2)(u_1, v_2).R_{\mathcal{C}(x_1, x_2)}. \end{aligned}$$

By Proposition 4.3, if $R_{\mathcal{A}}(B_1 \times B_2)(u_1, v_2).R_{\mathcal{C}(x_1, x_2)} \subset [u_1, v_2]_{\mathcal{A}, \mathcal{C}(x_1, x_2)}/R_{\mathcal{C}(x_1, x_2)}$, it would follow that $R_{\mathcal{A}}(B_1 \times B_2)(v_1, v_2).R_{\mathcal{C}} \subset [v_1, v_2]_{\mathcal{A}, \mathcal{C}}/R_{\mathcal{C}}$. Consider the sequence of fibrations in (4.16). Since

$$R_{\mathcal{A}}(B_1 \times B_2)(e, v_2).R_{\mathcal{C}^{(\infty)}} = R_{\mathcal{A}}(e, v_2(B_2 \cap M_{C_2(v_1, v_2)})).R_{\mathcal{C}^{(\infty)}} = \mathcal{Z}^{(\infty)},$$

we see inductively that $R_{\mathcal{A}}(B_1 \times B_2)(u_1^{(i)}, v_2).R_{\mathcal{C}^{(i)}} \subset \mathcal{Z}^{(i)}$ holds for all $i \geq 0$. $\square$

Proposition 4.10. *Given admissible triples $\mathcal{A} = (A_1, A_2, a, K)$ and $\mathcal{C} = (C_1, C_2, c, L)$, suppose that $\mathcal{A}' = (A_1, A_2, a, K')$ and $\mathcal{C}' = (C_1, C_2, c, L')$ are two other admissible quadruples containing the same triples (A_1, A_2, a) and (C_1, C_2, c) . Then there exist $t_2, s_2 \in T_2$ such that*

$$[v_1, v_2]_{\mathcal{A}', \mathcal{C}'} = (e, t_2)[v_1, v_2]_{\mathcal{A}, \mathcal{C}}(e, s_2), \quad \forall v_1 \in W_1^{C_1}, v_2 \in {}^{A_2}W_2.$$

Proof. Let K be given as in (2.1), and assume that

$$K' = \{(m_1, m_2) \in M_{A_1} \times M_{A_2} \mid \theta'_a(m_1 Z'_1) = m_2 Z'_2\},$$

where, for $i = 1, 2$, Z'_i is a closed subgroup of Z_{A_i} , and $\theta'_1: M_{A_1}/Z'_1 \rightarrow M_{A_2}/Z'_2$ is an isomorphism having the same properties as θ_a . The isomorphisms from M_{A_1}/Z_{A_1} to M_{A_2}/Z_{A_2} induced by θ_a and θ'_a will still be denoted by the same symbols. Then our assumptions imply that the automorphism $\theta := (\theta'_a)^{-1}\theta_a$ of M_{A_1}/Z_{A_1} is inner. Since θ leaves both T_1/Z_{A_1} and $(B_1 \cap M_{A_1})/Z_{A_1}$ invariant, $\theta = \text{Ad}_{t_1}$ for some $t_1 \in T_1$. It follows that $K'(B_1 \times B_2) = (e, t_2)K(B_1 \times B_2)$ for some $t_2 \in T_2$. Similarly, $(B_1 \times B_2)L' = (B_1 \times B_2)L(e, s_2)$ for some $s_2 \in T_2$. Proposition 4.10 now follows from Proposition 4.9. $\square$

Example 4.11. *Consider the case when $G_1 = G_2 = G$. Take $A_1 = A_2 = \Gamma$ and $a = \text{id}$, where Γ is the set of all simple roots for a pair (B, T) of Borel subgroup B of G and maximal torus $T \subset B$. Take $R_{\mathcal{A}} = K = G_{\text{diag}}$ and for some $t \in T$, take $R_{\mathcal{A}'} = K' = \{(g, \text{tgt}^{-1}) : g \in G\}$. Let $R_{\mathcal{C}} = R_{\mathcal{C}'} = B \times B$. Then it is easy to see that*

$$R_{\mathcal{A}'}(v, 1)(B \times B) = (e, t)R_{\mathcal{A}}(v, 1)(B \times B)$$

for all v in the Weyl group of (G, T) .

5. CLOSURES OF THE SETS $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}$ IN $G_1 \times G_2$

5.1. The set $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}$ for any $v_1 \in W_1$ and $v_2 \in W_2$. Let $\mathcal{A} = (A_1, A_2, a, K)$ and $\mathcal{C} = (C_1, C_2, c, L)$ be two arbitrary admissible quadruples for $G_1 \times G_2$. Extending (4.19), let

$$(5.1) \quad [v_1, v_2]_{\mathcal{A}, \mathcal{C}} = R_{\mathcal{A}}(B_1 \times B_2)(v_1, v_2)(B_1 \times B_2)R_{\mathcal{C}} \subset G_1 \times G_2, \quad \forall v_1 \in W_1, v_2 \in W_2.$$

Lemma 5.1. *When $v_1 \in W_1^{C_1}$,*

$$(5.2) \quad (B_1 \times B_2)(v_1, v_2)(B_1 \times B_2)R_{\mathcal{C}} = (B_1 \times B_2)(v_1, v_2)R_{\mathcal{C}}, \quad \forall v_2 \in W_2.$$

Proof. Since $v_1 \in W_1^{C_1}$, one has $B_1 v_1 (B_1 \cap M_{C_1}) = B_1 v_1$, and thus

$$\begin{aligned} (B_1 \times B_2)(v_1, v_2)(B_1 \times B_2)R_{\mathcal{C}} &= (B_1 \times B_2)(v_1, v_2)(B_1 \cap M_{C_1} \times \{e\})R_{\mathcal{C}} \\ &= (B_1 \times B_2)(v_1, v_2)R_{\mathcal{C}}.X_{(v_1, v_2)} \end{aligned}$$

□

When $v_1 \in W_1^{C_1}$ and $v_2 \in {}^{A_2}W_2$, the set $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}$ in (5.1) is thus the same as what we defined before. Our main result in this section is the following Theorem 5.2 which describes the closure of $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}$ in $G_1 \times G_2$ for any $v_1 \in W_1, v_2 \in W_2$. The proof of Theorem 5.2, which uses a series of lemmas proved in §5.2, will be given in §5.3. In this section, if X is a subset of $G_1 \times G_2$, $\overline{X}$ always denotes the closure of X in $G_1 \times G_2$.

Theorem 5.2. *For any $v_1 \in W_1, v_2 \in W_2$,*

$$(5.3) \quad \overline{[v_1, v_2]_{\mathcal{A}, \mathcal{C}}} = \bigsqcup_{\substack{v'_1 \in W_1^{C_1}, v'_2 \in {}^{A_2}W_2 : \\ \exists x_1 \in W_{A_1}, y_1 \in W_{C_1} \text{ s.t.} \\ x_1 v'_1 y_1 \leq v_1 \\ a(x_1) v'_2 c(y_1) \leq v_2}} [v'_1, v'_2]_{\mathcal{A}, \mathcal{C}} \quad (\text{disjoint union}).$$

Remark 5.3. A special case of (2.6) says that for any admissible quadruple $\mathcal{C}$, the $(B_1 \times B_2)$ -orbits in $(G_1 \times G_2)/R_{\mathcal{C}}$ are precisely of the form $(B_1 \times B_2)(v_1, v_2).R_{\mathcal{C}}$, where $v_1 \in W_1^{C_1}$ and $v_2 \in W_2$. Combining Lemma 5.1 and Theorem 5.2, we see that for any $(B_1 \times B_2)$ -orbit $\mathcal{O}$ in $(G_1 \times G_2)/R_{\mathcal{C}}$, the closure of the set $R_{\mathcal{A}}\mathcal{O}$ in $(G_1 \times G_2)/R_{\mathcal{C}}$ is a disjoint union of some sets of the form $R_{\mathcal{A}}(B_1 \times B_2)(v_1, v_2).R_{\mathcal{C}}$ with $v_1 \in W_1^{C_1}$ and $v_2 \in {}^{A_2}W_2$.

5.2. The closures of $(B_1 \times B_2, R_{\mathcal{C}})$ -double cosets in $G_1 \times G_2$. In this section, we will describe the $(B_1 \times B_2, R_{\mathcal{C}})$ -double cosets in the set

$$\overline{(B_1 \times B_2)(v_1, v_2)(B_1 \times B_2)R_{\mathcal{C}}} \subset G_1 \times G_2$$

for any $v_1 \in W_1, v_2 \in W_2$. Note again that by Lemma 5.1, $(B_1 \times B_2)(v_1, v_2)(B_1 \times B_2)R_{\mathcal{C}}$ is a single $(B_1 \times B_2, R_{\mathcal{C}})$ -double coset when $v_1 \in W_1^{C_1}$.

For $y_1, z_1 \in W_1$, let

$$\mathcal{W}_1(y_1, z_1) = \{x_1 \in W_1 \mid B_1 x_1 B_1 \subset B_1 y_1 B_1 z_1 B_1\}.$$

The following Lemma 5.4 on $\mathcal{W}_1(x_1, z_1)$ can be either proved by induction on the length of z_1 or can be seen as a direct consequence of the explicit description of $\mathcal{W}_1(y_1, z_1)$ in [2, Remark 3.19].

Lemma 5.4. *Let $y_1, z_1 \in W_1$ be arbitrary. Then every $x_1 \in \mathcal{W}_1(y_1, z_1)$ is of the form $x_1 = y_1 u_1$ for some $u_1 \in W_1, u_1 \leq z_1$.*

Lemma 5.5. *Let $y_1 \in W_1, y_2 \in W_2$. Then every $(B_1 \times B_2, R_{\mathcal{C}})$ -double coset in*

$$(B_1 \times B_2)(y_1, y_2)(B_1 \times B_2)R_{\mathcal{C}}$$

is of the form $(B_1 \times B_2)(y_1, u_2)R_{\mathcal{C}}$ for some $u_2 \in W_2, u_2 \leq y_2$.

Proof. Write $y_2 = w_2 z_2$, where $w_2 \in W_2^{C_2}$, $z_2 \in W_{C_2}$. Then

$$\begin{aligned}
(B_1 \times B_2)(y_1, y_2)(B_1 \times B_2)R_C &= ((B_1 y_1 B_1) \times (B_2 w_2 z_2))R_C \\
&= ((B_1 y_1 B_1 c^{-1}(z_2^{-1})) \times (B_2 w_2))R_C \\
&= ((B_1 y_1 B_1 c^{-1}(z_2^{-1})) \times (B_2 w_2 B_2))R_C \\
&= ((B_1 y_1 B_1 c^{-1}(z_2^{-1}) B_1) \times (B_2 w_2))R_C \\
&= \bigcup_{x_1 \in \mathcal{W}_1(y_1, c^{-1}(z_2^{-1}))} ((B_1 x_1 B_1) \times (B_2 w_2))R_C \\
&= \bigcup_{x_1 \in \mathcal{W}_1(y_1, c^{-1}(z_2^{-1}))} (B_1 \times B_2)(x_1, w_2)R_C.
\end{aligned}$$

By Lemma 5.4, every $x_1 \in \mathcal{W}_1(y_1, c^{-1}(z_2^{-1}))$ is of the form $x_1 = y_1 u_1$ for some $u_1 \in W_1$ such that $u_1 \leq c^{-1}(z_2^{-1})$, i.e. $c(u_1^{-1}) \leq z_2$. For such an $x_1 \in \mathcal{W}_1(y_1, c^{-1}(z_2^{-1}))$,

$$(B_1 \times B_2)(x_1, w_2)R_C = (B_1 \times B_2)(y_1, w_2 c(u_1^{-1}))R_C,$$

and $w_2 c(u_1^{-1}) \leq w_2 z_2 = y_2$. This completes the proof the Lemma. $\square$

Lemma 5.6. *For any $v_1 \in W_1$ and $v_2 \in W_2$, one has*

$$\overline{(B_1 \times B_2)(v_1, v_2)(B_1 \times B_2)R_C} = \overline{(B_1 \times B_2)(v_1, v_2)(B_1 \times B_2)} R_C.$$

Proof. The Lemma follows from [21, Lemma 2, P. 68] (see Lemma 8.1 in the Appendix) by noting that $R_C/((B_1 \times B_2) \cap R_C)$ is isomorphic to the full flag variety of M_{C_1} (see Lemma 4.5) and is hence complete. $\square$

Lemma 5.7. *For any $v_1 \in W_1$ and $v_2 \in W_2$, one has*

$$\begin{aligned}
(5.4) \quad \overline{(B_1 \times B_2)(v_1, v_2)(B_1 \times B_2)R_C} &= \bigcup_{\substack{w_1 \in W_1, w_2 \in W_2 : \\ w_1 \leq v_1, w_2 \leq v_2}} (B_1 \times B_2)(w_1, w_2)R_C.
\end{aligned}$$

Proof. By Lemma 5.6 and the Bruhat decomposition,

$$\begin{aligned}
\overline{(B_1 \times B_2)(v_1, v_2)(B_1 \times B_2)R_C} &= \bigcup_{\substack{y_1 \in W_1, y_2 \in W_2 : \\ y_1 \leq v_1, y_2 \leq v_2}} (B_1 \times B_2)(y_1, y_2)(B_1 \times B_2)R_C.
\end{aligned}$$

Let $y_1 \in W_1, y_2 \in W_2$ be such that $y_1 \leq v_1, y_2 \leq v_2$. By Lemma 5.5, every $(B_1 \times B_2, R_C)$ -double coset in $(B_1 \times B_2)(y_1, y_2)(B_1 \times B_2)R_C$ is of the form $(B_1 \times B_2)(y_1, u_2)R_C$ with $u_2 \in W_2, u_2 \leq y_2 \leq v_2$. Thus $\overline{(B_1 \times B_2)(v_1, v_2)(B_1 \times B_2)R_C}$ is contained in the right hand side of (5.4). Conversely, let $w_1 \in W_1, w_2 \in W_2$ be such that $w_1 \leq v_1, w_2 \leq v_2$. Then

$$\begin{aligned}
(B_1 \times B_2)(w_1, w_2)R_C &\subset (B_1 \times B_2)(w_1, w_2)(B_1 \times B_2)R_C \\
&\subset \overline{(B_1 \times B_2)(v_1, v_2)(B_1 \times B_2)} R_C.
\end{aligned}$$

Thus the right hand side of (5.4) is contained in $\overline{(B_1 \times B_2)(v_1, v_2)(B_1 \times B_2)R_C}$. $\square$

Proposition 5.8. *For any $v_1 \in W_1$ and $v_2 \in W_2$, one has the disjoint union*

$$\begin{aligned}
\overline{(B_1 \times B_2)(v_1, v_2)(B_1 \times B_2)R_C} &= \bigsqcup_{\substack{w_1 \in W_1^{C_1}, w_2 \in W_2 : \\ \exists u_1 \in W_{C_1} \text{ s.t.} \\ w_1 u_1 \leq v_1, w_2 c(u_1) \leq v_2}} (B_1 \times B_2)(w_1, w_2)R_C.
\end{aligned}$$

Proof. Proposition 5.8 follows from Lemma 5.7 by decomposing the element $w_1 \in W_1$ in the right hand side of (5.4) according to the decomposition $W_1 = W_1^{C_1} W_{C_1}$ and by the fact that $(T_1 \times T_2)(u_1, c(u_1))R_C = (T_1 \times T_2)R_C$ for any $u_1 \in W_{C_1}$, where T_1 and T_2 are respectively the maximal tori of G_1 and G_2 as fixed in §2.1. The union is disjoint by Theorem 3.1. $\square$

Corollary 5.9. *For any $v_1 \in W_1^{C_1}, v_2 \in W_2$, one has the disjoint union*

$$\overline{(B_1 \times B_2)(v_1, v_2)R_C} = \bigsqcup_{\substack{w_1 \in W_1^{C_1}, w_2 \in W_2 : \\ \exists u_1 \in W_{C_1} \text{ s.t.} \\ w_1 u_1 \leq v_1, w_2 c(u_1) \leq v_2}} (B_1 \times B_2)(w_1, w_2)R_C.$$

Remark 5.10. By Lemma 8.3 in the Appendix, Corollary 5.9 is equivalent to the following description of closures of $(B_1 \times B_2, R_C^-)$ -double cosets in $G_1 \times G_2$ as given in [19, Lemma 2.2]: for $v_1 \in W_1^{C_1}, v_2 \in W_2$,

$$\overline{(B_1 \times B_2)(v_1, v_2)R_C^-} = \bigsqcup_{\substack{w_1 \in W_1^{C_1}, w_2 \in W_2 : \\ \exists u_1 \in W_{C_1} \text{ s.t.} \\ v_1 u_1^{-1} \leq w_1, w_2 c(u_1) \leq v_2}} (B_1 \times B_2)(w_1, w_2)R_C^-.$$

Lemma 5.11. *Let $w_1 \in W_1, w_2 \in W_2, v'_1 \in W_1^{C_1}$, and $v'_2 \in {}^{A_2}W_2$. Suppose that*

$$(5.5) \quad (R_A(B_1 \times B_2)(w_1, w_2)R_C) \cap [v'_1, v'_2]_{A,C} \neq \emptyset.$$

Then there exist $x_1 \in W_{A_1}$ and $y_1 \in W_{C_1}$ such that

$$x_1 v'_1 y_1 \leq w_1 \quad \text{and} \quad a(x_1) v'_2 c(y_1) \leq w_2.$$

Proof. One sees from (5.5) that $(w_1, w_2)R_C \subset (B_1 \times B_2)R_A(B_1 \times B_2)(v'_1, v'_2)(B_1 \times B_2)R_C$. Since

$$(B_1 \times B_2)R_A(B_1 \times B_2) = \bigcup_{x_1 \in W_{A_1}} (B_1 \times B_2)(x_1, a(x_1))(B_1 \times B_2),$$

we have

$$\begin{aligned} (w_1, w_2)R_C &\subset \bigcup_{x_1 \in W_{A_1}} ((B_1 x_1 B_1 v'_1 B_1) \times (B_2 a(x_1) B_2 v'_2 B_2)) R_C \\ &= \bigcup_{x_1 \in W_{A_1}} ((B_1 x_1 B_1 v'_1 B_1) \times (B_2 a(x_1) v'_2 B_2)) R_C, \end{aligned}$$

where in the last step we used the fact that $l(a(x_1)v'_2) = l(a(x_1)) + l(v'_2)$. Thus there exists $x_1 \in W_{A_1}$ such that $(w_1, w_2) \in ((B_1 x_1 B_1 v'_1 B_1) \times (B_2 a(x_1) v'_2 B_2))R_C$. Since

$$R_C \subset \bigcup_{y_1 \in W_{C_1}} ((B_1 y_1 B_1) \times (B_2 c(y_1) B_2)),$$

there exists $y_1 \in W_{C_1}$ such that

$$\begin{aligned} (w_1, w_2) &\in (B_1 x_1 B_1 v'_1 B_1 y_1 B_1) \times (B_2 a(x_1) v'_2 B_2 c(y_1) B_2) \\ &= (B_1 x_1 B_1 v'_1 y_1 B_1) \times (B_2 a(x_1) v'_2 B_2 c(y_1) B_2). \end{aligned}$$

By Lemma 8.4 in the Appendix, $w_1 \geq x_1 v'_1 y_1$ and $w_2 \geq a(x_1) v'_2 c(y_1)$. $\square$

5.3. Proof of Theorem 5.2. Fix $v_1 \in W_1$ and $v_2 \in W_2$. Let

$$\mathcal{J}(v_1, v_2) = \{(v'_1, v'_2) \in W_1^{C_1} \times {}^{A_2}W_2 \mid \overline{[v_1, v_2]_{\mathcal{A}, \mathcal{C}}} \cap [v'_1, v'_2]_{\mathcal{A}, \mathcal{C}} \neq \emptyset\}.$$

Then

$$\overline{[v_1, v_2]_{\mathcal{A}, \mathcal{C}}} = \bigsqcup_{(v'_1, v'_2) \in \mathcal{J}(v_1, v_2)} \overline{[v_1, v_2]_{\mathcal{A}, \mathcal{C}}} \cap [v'_1, v'_2]_{\mathcal{A}, \mathcal{C}}.$$

We will first show that

$$(5.6) \quad \mathcal{J}(v_1, v_2) = \{(v'_1, v'_2) \in W_1^{C_1} \times {}^{A_2}W_2 \mid \\ \exists x_1 \in W_{A_1}, y_1 \in W_{C_1} \text{ s.t. } x_1 v'_1 y_1 \leq v_1, a(x_1) v'_2 c(y_1) \leq v_2\}.$$

Indeed, by Lemma 8.1 in the Appendix,

$$\overline{[v_1, v_2]_{\mathcal{A}, \mathcal{C}}} = R_{\mathcal{A}} \overline{(B_1 \times B_2)(v_1, v_2)(B_1 \times B_2)} R_{\mathcal{C}}.$$

Thus by Lemma 5.7,

$$\overline{[v_1, v_2]_{\mathcal{A}, \mathcal{C}}} = \bigsqcup_{\substack{w_1 \in W_1, w_2 \in W_2 : \\ w_1 \leq v_1, w_2 \leq v_2}} R_{\mathcal{A}}(B_1 \times B_2)(w_1, w_2) R_{\mathcal{C}}.$$

Suppose that $(v'_1, v'_2) \in \mathcal{J}(v_1, v_2)$. Then there exists $(w_1, w_2) \in W_1 \times W_2$ with $w_1 \leq v_1, w_2 \leq v_2$ such that

$$(R_{\mathcal{A}}(B_1 \times B_2)(w_1, w_2) R_{\mathcal{C}}) \cap [v'_1, v'_2]_{\mathcal{A}, \mathcal{C}} \neq \emptyset.$$

By Lemma 5.11, there exist $x_1 \in W_{A_1}, y_1 \in W_{C_1}$ such that

$$x_1 v'_1 y_1 \leq w_1 \leq v_1 \quad \text{and} \quad a(x_1) v'_2 c(y_1) \leq w_2 \leq v_2.$$

Thus (v'_1, v'_2) is in the set of the right hand side of (5.6). Conversely, suppose that $(v'_1, v'_2) \in W_1^{C_1} \times {}^{A_2}W_2$ are such that $x_1 v'_1 y_1 \leq v_1$ and $a(x_1) v'_2 c(y_1) \leq v_2$ for some $x_1 \in W_{A_1}$ and $y_1 \in W_{C_1}$. Let $w_1 = x_1 v'_1 y_1$ and $w_2 = a(x_1) v'_2 c(y_1)$ so that

$$v'_1 = x_1^{-1} w_1 y_1^{-1} \quad \text{and} \quad v'_2 = a(x_1^{-1}) w_2 c(y_1^{-1}).$$

It follows that $(v'_1 T_1, v'_2 T_2) = (x_1^{-1} w_1 y_1^{-1} T_1, a(x_1^{-1}) w_2 c(y_1^{-1}) T_2)$ and hence

$$(5.7) \quad (v'_1 T_1, v'_2 T_2) \subset R_{\mathcal{A}}(B_1 \times B_2)(w_1, w_2) R_{\mathcal{C}} \subset \overline{[v_1, v_2]_{\mathcal{A}, \mathcal{C}}},$$

where $T_i = B_i^- \cap B_i$ for $i = 1, 2$. Thus $\overline{[v_1, v_2]_{\mathcal{A}, \mathcal{C}}} \cap [v'_1, v'_2]_{\mathcal{A}, \mathcal{C}} \neq \emptyset$, and so $(v'_1, v'_2) \in \mathcal{J}(v_1, v_2)$. $\square$

For any $(v'_1, v'_2) \in \mathcal{J}(v_1, v_2)$, it follows from (5.7) that

$$R_{\mathcal{A}}(v'_1, v'_2 T_2) R_{\mathcal{C}} = R_{\mathcal{A}}(v'_1 T_1, v'_2 T_2) R_{\mathcal{C}} \subset \overline{[v_1, v_2]_{\mathcal{A}, \mathcal{C}}}.$$

By Lemma 3.3, $R_{\mathcal{A}}(v'_1, v'_2 T_2) R_{\mathcal{C}}$ is dense in $[v'_1, v'_2]_{\mathcal{A}, \mathcal{C}}$. Hence $[v'_1, v'_2]_{\mathcal{A}, \mathcal{C}} \subset \overline{[v_1, v_2]_{\mathcal{A}, \mathcal{C}}}$, and $\overline{[v_1, v_2]_{\mathcal{A}, \mathcal{C}}} \cap [v'_1, v'_2]_{\mathcal{A}, \mathcal{C}} = [v'_1, v'_2]_{\mathcal{A}, \mathcal{C}}$. This completes the proof of Theorem 5.2.

6. THE $(R_{\mathcal{A}}, R_{\mathcal{C}}^-)$ -STABLE SUBSETS $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}^-$

6.1. The subsets $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}^-$ of $G_1 \times G_2$. Let again $\mathcal{A}$ and $\mathcal{C}$ be two admissible quadruples for $G_1 \times G_2$. Let w_{0, Γ_1} and w_{0, C_1} be the longest element in W_1 and W_{C_1} respectively.

Associated to $\mathcal{C}$, we have the admissible quadruple $\mathcal{C}^* \stackrel{\text{def}}{=} (C_1^*, C_2^*, c^*, L^*)$, where

$$(6.1) \quad C_1^* = -w_{0, \Gamma_1}(C_1), \quad C_2^* = C_2, \quad c^* = c(w_{0, \Gamma_1} w_{0, C_1})^{-1}, \quad L^* = \text{Ad}_{(\dot{w}_{0, \Gamma_1} \dot{w}_{0, C_1}, e)} L.$$

It is easy to see that $R_{\mathcal{C}^*} = \text{Ad}_{(\dot{w}_{0, \Gamma_1} \dot{w}_{0, C_1}, e)} R_{\mathcal{C}}^-$. Moreover,

$$(6.2) \quad W_1^{C_1}(w_{0, \Gamma_1} w_{0, C_1})^{-1} = W_1^{C_1^*}$$

Proposition 6.1. For $v_1 \in W_1^{C_1}$ and $v_2 \in {}^{A_2}W_2$, let

$$[v_1, v_2]_{\mathcal{A}, \mathcal{C}}^- = R_{\mathcal{A}}(v_1, v_2 M_{C_2(v_1, v_2)}) R_{\mathcal{C}}^- \subset G_1 \times G_2.$$

Then the following properties hold:

(i)

$$G_1 \times G_2 = \bigsqcup_{v_1 \in W_1^{C_1}, v_2 \in {}^{A_2}W_2} [v_1, v_2]_{\mathcal{A}, \mathcal{C}}^- \quad (\text{disjoint union});$$

(ii) $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}^- = R_{\mathcal{A}}(B_1 \times B_2)(v_1, v_2) R_{\mathcal{C}}^-$ for every $v_1 \in W_1^{C_1}$ and $v_2 \in {}^{A_2}W_2$;

(iii) $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}^-$ is locally closed, smooth, and irreducible. Its projection $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}^- / R_{\mathcal{C}}^-$ to $(G_1 \times G_2) / R_{\mathcal{C}}^-$ fibers over the flag variety $M_{A_1} / (M_{A_1} \cap P_{A_1(v_1, v_2)})$ with fibers isomorphic to the product of $M_{C_2(v_1, v_2)} / Y_2$ and the affine space of dimension $\dim U_{A_1(v_1, v_2)} - l(v_1) + l(v_2)$, where Y_2 is as in Theorem 2.2.

Proof. Let $v_1^* = v_1(w_{0, \Gamma_1} w_{0, C_1})^{-1} \in W_1^{C_1^*}$. All the statements in Proposition 6.1 follow from the fact that

$$[v_1, v_2]_{\mathcal{A}, \mathcal{C}}^- = R_{\mathcal{A}}(B_1 \times B_2)(v_1^*, v_2)(B_1 \times B_2) R_{\mathcal{C}}^*(\dot{w}_{0, \Gamma_1} \dot{w}_{0, C_1}, e).$$

□

6.2. Closures of the sets $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}^-$ in $G_1 \times G_2$. For each $v_1 \in W_1$ and $v_2 \in W_2$, set

$$(6.3) \quad [v_1, v_2]_{\mathcal{A}, \mathcal{C}}^- = R_{\mathcal{A}}(B_1 \times B_2)(v_1, v_2)(w_{0, C_1}(B_1^-) \times B_2) R_{\mathcal{C}}^- \subset G_1 \times G_2.$$

It follows from Lemma 5.1 that

$$(6.4) \quad [v_1, v_2]_{\mathcal{A}, \mathcal{C}}^- = R_{\mathcal{A}}(B_1 \times B_2)(v_1, v_2) R_{\mathcal{C}}^-, \quad \text{when } v_1 \in W_1^{C_1}, v_2 \in W_2.$$

Theorem 6.2. For any $v_1 \in W_1$ and $v_2 \in W_2$, with $[v_1, v_2]_{\mathcal{A}, \mathcal{C}}^-$ given in (6.3), one has

$$\begin{aligned} \overline{[v_1, v_2]_{\mathcal{A}, \mathcal{C}}^-} &= \bigsqcup_{\substack{v'_1 \in W_1^{C_1}, v'_2 \in {}^{A_2}W_2 : \\ \exists x_1 \in W_{A_1}, y_1 \in W_{C_1} \text{ s.t.} \\ x_1 v'_1 y_1 w_{0, C_1} \geq v_1 w_{0, C_1} \\ a(x_1) v'_2 c(y_1) \leq v_2}} [v'_1, v'_2]_{\mathcal{A}, \mathcal{C}}^- \end{aligned}$$

Proof. For $v_1 \in W_1$, let again $v_1^* = v_1(w_{0, \Gamma_1} w_{0, C_1})^{-1}$. Then

$$[v_1, v_2]_{\mathcal{A}, \mathcal{C}}^- = R_{\mathcal{A}}(B_1 \times B_2)(v_1^*, v_2)(B_1 \times B_2) R_{\mathcal{C}}^*(\dot{w}_{0, \Gamma_1} \dot{w}_{0, C_1}, e), \quad \forall v_1 \in W_1, v_2 \in W_2.$$

For $y_1 \in W_{C_1}$, set $(y_1)_* = (w_{0, \Gamma_1} w_{0, C_1}) y_1 (w_{0, \Gamma_1} w_{0, C_1})^{-1} \in W_{C_1^*}$. By Theorem 5.2,

$$\begin{aligned} \overline{[v_1, v_2]_{\mathcal{A}, \mathcal{C}}^-} &= \bigsqcup_{\substack{v'_1 \in W_1^{C_1}, v'_2 \in {}^{A_2}W_2 : \\ \exists x_1 \in W_{A_1}, y_1 \in W_{C_1} \text{ s.t.} \\ x_1 (v'_1)^* (y_1)_* \leq v_1^* \\ a(x_1) v'_2 c(y_1) \leq v_2}} [v'_1, v'_2]_{\mathcal{A}, \mathcal{C}}^- \\ &= \bigsqcup_{\substack{v'_1 \in W_1^{C_1}, v'_2 \in {}^{A_2}W_2 : \\ \exists x_1 \in W_{A_1}, y_1 \in W_{C_1} \text{ s.t.} \\ x_1 v'_1 y_1 w_{0, C_1} \geq v_1 w_{0, C_1} \\ a(x_1) v'_2 c(y_1) \leq v_2}} [v'_1, v'_2]_{\mathcal{A}, \mathcal{C}}^- \end{aligned}$$

□

7. THE $R_{\mathcal{A}}$ -STABLE PIECES IN $\overline{G}$ AND THEIR CLOSURES

We retain the notation in §2.3. In particular, G is a semi-simple algebraic group of adjoint type, and $\overline{G}$ denotes the De Concini-Procesi compactification of G . In this section, unless otherwise stated, if X is a subset of $\overline{G}$, $\overline{X}$ always denotes the closure of X in $\overline{G}$.

For each $J \subset G$, recall that h_J is a point in $\overline{G}$ such that the stabilizer subgroup of $G \times G$ at h_J is R_J^- given in (2.8). Let $\mathcal{A} = (A_1, A_2, a, K)$ be any admissible quadruple for $G \times G$. Then we have the decomposition of $\overline{G}$ into $R_{\mathcal{A}}$ -stable pieces

$$\overline{G} = \bigsqcup_{J \subset \Gamma, v_1 \in W^J, v_2 \in {}^{A_2}W} [J, v_1, v_2]_{\mathcal{A}},$$

where for $J \subset \Gamma$ and for $v_1 \in W^J$ and $v_2 \in {}^{A_2}W$, the subset $[J, v_1, v_2]_{\mathcal{A}}$ of $\overline{G}$ is defined by (2.9), namely

$$[J, v_1, v_2]_{\mathcal{A}} = R_{\mathcal{A}}(B \times B)(v_1, v_2).h_J.$$

By part (iii) of Proposition 6.1, we have

Proposition 7.1. *Let $\mathcal{A}$ be any admissible quadruple for $G \times G$. Then for any $J \subset \Gamma$ and $v_1 \in W^J$, $v_2 \in {}^{A_2}W$, $[J, v_1, v_2]_{\mathcal{A}}$ is a locally closed smooth subset of $\overline{G}$. It fibers over the flag variety $M_{A_1}/(M_{A_1} \cap P_{A_1(v_1, v_2)})$ with fibers isomorphic to the product of $M_{J(v_1, v_2)}/Z_J$ and the affine space of dimension $\dim U_{A_1(v_1, v_2)} - l(v_1) + l(v_2)$, where $J(v_1, v_2)$ is the smallest subset of J stable under $v_2^{-1}av_1$ and $A_1(v_1, v_2) = v_1 J(v_1, v_2) \subset A_1$.*

When $R_{\mathcal{A}} = G_{\text{diag}}$, the Proposition 7.1 coincides with the description of the geometry of Lusztig's G_{diag} -stable pieces given in [7].

In this section, we study the closures of the subsets $[J, v_1, v_2]_{\mathcal{A}}$ in $\overline{G}$.

7.1. The first description. For $J \subset \Gamma$ and $v_1 \in W^J$, $v_2 \in W$, let

$$[J, v_1, v_2] = (B \times B)(v_1, v_2).h_J \subset \overline{G}.$$

The following Lemma 7.2 follows immediately from Lemma 8.1 in the Appendix.

Lemma 7.2. *For any admissible triple $\mathcal{A}$ for $G \times G$ and all $J \subset \Gamma$, $v_1 \in W^J$, $v_2 \in {}^{A_2}W$,*

$$\overline{[J, v_1, v_2]_{\mathcal{A}}} = R_{\mathcal{A}}\overline{[J, v_1, v_2]}.$$

Proposition 7.3. *Let $\mathcal{A}$ be an admissible quadruple for $G \times G$. Then for any $J \subset \Gamma$ and $(v_1, v_2) \in W^J \times {}^{A_2}W$,*

$$(7.1) \quad \overline{[J, v_1, v_2]_{\mathcal{A}}} = \bigsqcup_{I \subset J} \bigsqcup_{\substack{v'_1 \in W^I, v'_2 \in {}^{A_2}W : \\ \exists x \in W_{A_1}, y \in W_I, z \in W_J^I \text{ s.t.} \\ l(v_2 z) = l(v_2) + l(z) \\ xv'_1 y w_{0,I} \geq v_1 z w_{0,I} \\ a(x)v'_2 y \leq v_2 z}} [I, v'_1, v'_2]_{\mathcal{A}}.$$

Proof. It is well-known [4] that for $I_1, I_2 \subset \Gamma$,

$$\overline{(G \times G).h_{I_1}} \cap (G \times G).h_{I_2} \neq \emptyset$$

if and only if $I_2 \subset I_1$. Thus

$$\overline{[J, v_1, v_2]_{\mathcal{A}}} = \bigsqcup_{I \subset J} \left(\overline{[J, v_1, v_2]_{\mathcal{A}}} \cap (G \times G).h_I \right).$$

Fix $I \subset J$. By Lemma 7.2,

$$\overline{[J, v_1, v_2]_{\mathcal{A}}} \cap (G \times G).h_I = R_{\mathcal{A}} \left(\overline{[J, v_1, v_2]} \cap (G \times G).h_I \right).$$

By [19, Lemma 2.3],

$$\overline{[J, v_1, v_2]} \cap (G \times G).h_I = \bigsqcup_{\substack{z \in W_J^I, \\ l(v_2 z) = l(v_2) + l(z)}} \overline{[I, v_1 z, v_2 z]}^I,$$

where, for a subsets Y of $(G \times G).h_I$, $\overline{Y}^I$ denotes the closure of Y in $(G \times G).h_I$. Thus

$$\overline{[J, v_1, v_2]}_{\mathcal{A}} \cap (G \times G).h_I = \bigsqcup_{\substack{z \in W_J^I, \\ l(v_2 z) = l(v_2) + l(z)}} R_{\mathcal{A}} \overline{[I, v_1 z, v_2 z]}^I.$$

By Lemma 8.1, $R_{\mathcal{A}} \overline{[I, v_1 z, v_2 z]}^I = \overline{R_{\mathcal{A}}[I, v_1 z, v_2 z]}^I$ for every $z \in W_J^I$. The decomposition in (7.1) now follows from Theorem 6.2. $\square$

In the following §7.2 and §7.3, we will simplify the descriptions of the the closure relations in Proposition 7.3.

7.2. The second description. Recall that $(G \times G).h_{\emptyset} \cong (G \times G)/(B_- \times B)$ is the unique closed $(G \times G)$ -orbit in $\overline{G}$. For $J \subset \Gamma$ and $v_1 \in W_J$, $v_2 \in {}^{A_2}W$, let

$$\partial_{\emptyset} \overline{[J, v_1, v_2]}_{\mathcal{A}} = \overline{[J, v_1, v_2]}_{\mathcal{A}} \cap (G \times G).h_{\emptyset}.$$

Theorem 7.4. *Let $\mathcal{A}$ be an admissible quadruple for $G \times G$. Let $I, J \subset \Gamma$, $v_1 \in W_J$, $v'_1 \in W^I$, and $v_2, v'_2 \in {}^{A_2}W$. Then the following are equivalent:*

- (i) $[I, v'_1, v'_2]_{\mathcal{A}} \subset \overline{[J, v_1, v_2]}_{\mathcal{A}}$;
- (ii) $I \subset J$ and $\partial_{\emptyset} \overline{[I, v'_1, v'_2]}_{\mathcal{A}} \subset \partial_{\emptyset} \overline{[J, v_1, v_2]}_{\mathcal{A}}$;
- (iii) $I \subset J$ and $[\emptyset, v'_1, v'_2]_{\mathcal{A}} \subset \partial_{\emptyset} \overline{[J, v_1, v_2]}_{\mathcal{A}}$.

Proof. Clearly (i) implies (ii). Since $[\emptyset, v'_1, v'_2]_{\mathcal{A}} \subset \partial_{\emptyset} \overline{[I, v'_1, v'_2]}_{\mathcal{A}}$, (ii) implies (iii). It remains to show that (iii) implies (i).

Assume (iii). By Proposition 7.3, there exist $x \in W_{A_1}$ and $z \in W_J$ with $l(v_2 z) = l(v_2) + l(z)$ such that $xv'_1 \geq v_1 z$ and $a(x)v'_2 \leq v_2 z$. Write $z = uy$ with $u \in W_J^I$ and $y \in W_I$. Then the set

$$S = \{(x', y') \in W_{A_1} \times W_I \mid x'v'_1 \geq v_1 uy', \ a(x')v'_2 \leq v_2 uy'\}$$

is non-empty. Let $(x_0, y_0) \in W_{A_1} \times W_I$ be a minimal element in S . We claim that

$$(7.2) \quad (x_0 v'_1)^I \geq v_1 u y_0 (x_0 v'_1)^{-1}_I \quad \text{and} \quad a(x_0) v'_2 y_0^{-1} \leq v_2 u,$$

where $(x_0 v'_1)^I \in W^I$ and $(x_0 v'_1)_I \in W_I$ are such that $x_0 v'_1 = (x_0 v'_1)^I (x_0 v'_1)_I$. By Lemma 8.3, it would follow from (7.2) that

$$x_0 v'_1 y_0^{-1} w_{0,I} \geq v_1 u w_{0,I} \quad \text{and} \quad a(x_0) v'_2 y_0^{-1} \leq v_2 u,$$

and, since $l(v_2 u) = l(v_2) + l(u)$, we would see by Proposition 7.3 that $[I, v'_1, v'_2]_{\mathcal{A}} \subset \overline{[J, v_1, v_2]}_{\mathcal{A}}$.

It remains to prove (7.2). We first show that $a(x_0) v'_2 y_0^{-1} \leq v_2 u$. Indeed, since $a(x_0) v'_2 \leq v_2 u y_0$, it follows from Lemma 8.7 in the Appendix that there exists $y_1 \leq y_0$ such that

$$(7.3) \quad v_2 u = v_2 u y_0 y_0^{-1} \geq a(x_0) v'_2 y_1^{-1}.$$

Again by Lemma 8.7, there exists $y_2 \leq y_1$ such that

$$a(x_0) v'_2 = a(x_0) v'_2 y_1^{-1} y_1 \leq v_2 u y_2.$$

Since $y_2 \leq y_1 \leq y_0$, we have $x_0 v'_1 \geq v_1 u y_0 \geq v_1 u y_2$. Thus $(x_0, y_2) \in S$. Since $(x_0, y_2) \leq (x_0, y_0)$ and since (x_0, y_0) is minimal in S , we must have $y_2 = y_0$. Hence $y_1 = y_0$, and $a(x_0) v'_2 y_0^{-1} \leq v_2 u$ by (7.3).

We now show that $l(x_0v'_1) = l(x_0) + l(v'_1)$. If $l(x_0v'_1) < l(x_0) + l(v'_1)$, then by Lemma 8.6 in the Appendix, there exists $x_1 < x_0$ such that $x_1v'_1 > x_0v'_1 \geq v_1uy_0$. Since $a(x_1)v'_2 \leq a(x_0)v'_2 \leq v_2uy_0$, we have $(x_1, y_0) \in S$. Since $(x_1, y_0) < (x_0, y_0)$, this is a contradiction to the minimality of (x_0, y_0) in S . Hence $l(x_0v'_1) = l(x_0) + l(v'_1)$.

By Lemma 8.7 in the Appendix, there exists $y_1 \leq (x_1v'_1)_I$ such that

$$(x_0v'_1)^I = (x_0v'_1)(x_0v'_1)_I^{-1} \geq v_1uy_0y_1^{-1}.$$

By Lemma 8.8 in the Appendix, there exists $x_2 \leq x_0$ such that $(x_0v'_1)^I y_1 = x_2v'_1$. Now

$$x_2v'_1 = (x_0v'_1)^I y_1 \geq v_1uy_0y_1^{-1}y_1 = v_1uy_0,$$

and $a(x_2)v'_2 \leq a(x_0)v'_2 \leq v_2uy_0$. Hence $(x_2, y_0) \in S$. By the minimality of (x_0, y_0) in S , we have $x_2 = x_0$, so $y_1 = (x_1v'_1)_I$, and $(x_0v'_1)^I \geq v_1uy_0(x_1v'_1)_I^{-1}$. This proves (7.2). $\square$

As a corollary of (iii) in Theorem 7.4 and Proposition 7.3, we get the following second description of the closure relations on the $R_{\mathcal{A}}$ -stable pieces in $\overline{G}$.

Corollary 7.5. *Let $\mathcal{A}$ be an admissible quadruple for $G \times G$. Then for any $J \subset \Gamma$ and $(v_1, v_2) \in W^J \times {}^{A_2}W$,*

$$(7.4) \quad \overline{[J, v_1, v_2]_{\mathcal{A}}} = \bigsqcup_{I \subset J} \bigsqcup_{\substack{v'_1 \in W^I, v'_2 \in {}^{A_2}W : \\ \exists x \in W_{A_1}, z \in W_J \text{ s.t.} \\ l(v_2z) = l(v_2) + l(z) \\ xv'_1 \geq v_1z, \quad a(x)v'_2 \leq v_2z}} [I, v'_1, v'_2]_{\mathcal{A}}.$$

7.3. The third description. In this section, we show that the condition $l(v_2z) = l(v_2) + l(z)$ in (7.4) can be dropped.

Theorem 7.6. *Let $\mathcal{A}$ be an admissible quadruple for $G \times G$. Then for any $J \subset \Gamma$ and $(v_1, v_2) \in W^J \times {}^{A_2}W$,*

$$(7.5) \quad \overline{[J, v_1, v_2]_{\mathcal{A}}} = \bigsqcup_{I \subset J} \bigsqcup_{\substack{v'_1 \in W^I, v'_2 \in {}^{A_2}W : \\ \exists x \in W_{A_1}, z \in W_J \text{ s.t.} \\ xv'_1 \geq v_1z, \quad a(x)v'_2 \leq v_2z}} [I, v'_1, v'_2]_{\mathcal{A}}.$$

Proof. Fix $J \subset \Gamma$ and $(v_1, v_2) \in W^J \times {}^{A_2}W$. It is enough to show that the right hand side of (7.5) is contained in the right hand side of (7.4). To this end, let $I \subset J$ and let $(v'_1, v'_2) \in W^I \times {}^{A_2}W$ be such that there exist $x \in W_{A_1}$ and $z \in W_J$ with $xv'_1 \geq v_1z$ and $a(x)v'_2 \leq v_2z$. Choose such an $x \in W_{A_1}$ and let

$$Z = \{z' \in W_J \mid xv'_1 \geq v_1z', \quad a(x)v'_2 \leq v_2z'\}.$$

Then $Z \neq \emptyset$. Let $z_0 \in Z$ be a minimal element. We claim that $l(v_2z_0) = l(v_2) + l(z_0)$. Indeed, if $l(v_2z_0) < l(v_2) + l(z_0)$, then by Lemma 8.6 in the Appendix, there exists $z_1 < z_0$ such that $v_2z_1 > v_2z_0$. Thus $v_2z_1 > a(x)v'_2$. Since $v_1 \in W^J$ and $z_1, z_0 \in W_J$, we also have $v_1z_1 < v_1z_0 \leq xv'_1$. Thus $z_1 \in Z$, which contradicts to the fact the z_0 is a minimal element in Z . This shows that $l(v_2z_0) = l(v_2) + l(z_0)$, and thus $[I, v'_1, v'_2]_{\mathcal{A}}$ is contained in the right hand side of (7.4). $\square$

Remark 7.7. Note that in the proofs of Proposition 7.3, Corollary 7.5, and Theorem 7.6, we did not use the fact that $v_2 \in {}^{A_2}W$. In fact, the decomposition formulas (7.1), (7.4), and (7.5) hold for any $v_1 \in W^J$ and $v_2 \in W_2$. Thus the closure of $R_{\mathcal{A}}\mathcal{O}$ in $\overline{G}$ for any $(B \times B)$ -orbit $\mathcal{O}$ in $\overline{G}$ is a union of the sets of the form $[I, v'_1, v'_2]_{\mathcal{A}}$ for $I \subset \Gamma$, $v'_1 \in W^I$, and $v'_2 \in {}^{A_2}W$. Proposition 7.3, Corollary 7.5, and Theorem 7.6 give three equivalent descriptions of the decomposition. See also Remark 5.3.

8. APPENDIX

8.1. A lemma from [21]. The following is [21, Lemma 2, P. 68]. We state it here for the convenience of the reader.

Lemma 8.1. *Let G be an algebraic group acting on a variety V . Let H be a closed subgroup of G and let $U \subset V$ be a closed subset of V , invariant under the action of H . Assume that G/H is complete. Then $G.U$ is closed.*

8.2. A few facts on Weyl groups. Let G be a any connected reductive algebraic group over an algebraically closed field. Let T be a maximal torus of G , let B a Borel subgroup of G containing T , and let Γ be the set of simple roots for (B, T) . Let W be the Weyl group of Γ . For the convenience of the reader, we collect in this section a few facts on W that are used in this paper. Recall that for two subsets A and C of Γ we denote ${}^AW^C = {}^AW \cap W^C$.

Lemma 8.2. [3, Proposition 2.7.5] *For any $A, C \subset \Gamma$, every $v \in {}^AW$ can be uniquely written as a product $v = xu$, where $x \in {}^AW^C$ and $u \in {}^{C \cap x^{-1}(A)}W_C$.*

For $C \subset \Gamma$, let $w_{0,C}$ be the longest element of W_C .

Lemma 8.3. [19, Page 79] *Let $x, w \in W^C$ and $u \in W_C$. Then $xu^{-1} \leq w$ if and only if $xw_{0,C} \leq wuw_{0,C}$.*

Proof. If $xu^{-1} \leq w$, then $xw_{0,C} = xu^{-1}uw_{0,C} \leq wuw_{0,C}$. Conversely, assume that $xw_{0,C} \leq wuw_{0,C}$. Then there exist $w_1 \leq w$ and $y \leq uw_{0,C}$ such that $xw_{0,C} = w_1y$. It follows from $y \leq uw_{0,C}$ that $yw_{0,C} \geq u$, so $u^{-1} \leq w_{0,C}y^{-1}$. Thus $xu^{-1} \leq xw_{0,C}y^{-1} = w_1 \leq w$. $\square$

For $y, z \in W$, let $\mathcal{W}(y, z) = \{x \in W \mid BxB \subset ByBzB\}$.

Lemma 8.4. [18, p. 420] *Let $y, z \in W$. Then $x \geq yz$ for any $x \in \mathcal{W}(y, z)$.*

Lemma 8.5. [7, Lemma 3.3] *For any $u, w \in W$, the subset $\{vw \mid v \leq u\}$ of W contains a unique maximal element u_1w . Moreover, $l(u_1w) = l(u_1) + l(w)$.*

Lemma 8.6. *If $u, w \in W$ are such that $l(uw) < l(u) + l(w)$, then there exists u_1 such that $u_1 < u$ and $u_1w > uw$.*

Proof. The u_1 such that u_1w is the maximal element in the set $\{vw \mid v \leq u\}$ is as required. $\square$

Lemma 8.7. [7, Corollary 3.4] *Let $u, w, w' \in W$ and assume that $w' \leq w$. Then*

- (i) *there exists $u_1 \leq u$ such that $w'u_1 \leq wu$;*
- (ii) *there exists $u_2 \leq u$ such that $w'u \leq wu_2$.*

Lemma 8.8. [7, Lemma 3.10] *Let $J \subset \Gamma, w \in W^J$, and $u \in W$ be such that $l(uw) = l(u) + l(w)$. Write $uw = xv$ with $x \in W^J$ and $v \in W_J$. Then for any $v' \leq v$, there exists $u' \leq u$ such that $u'w = xv'$.*

REFERENCES

- [1] A. Belavin and V. Drinfeld, *Triangular equations and simple Lie algebras*, Math. Phys. Rev. **4** (1984), 93-165.
- [2] A. Borel and J. Tits, *Compléments à l'article: Groupes Réductifs*, Publ. Math. Inst. Hautes Etud. Sci. **41** (1972) 253-276.
- [3] R. W. Carter, *Finite groups of Lie type. Conjugacy classes and complex characters*. Wiley-Interscience Publ., Chichester, 1993.
- [4] C. De Concini and C. Procesi, *Complete symmetric varieties*, in Invariant theory (Montecatini, 1982), Lecture Notes in Math. **996** (1983), 1-44.
- [5] C. De Concini and T. Springer, *Compactification of symmetric spaces*, Transformation Groups **4** (1999), 273-300.
- [6] X. He, *Unipotent variety in the group compactification*, Adv. Math. **203** (2006), 109-131.

- [7] X. He, *The G -stable pieces of the wonderful compactification*, math.RT/0412302, to appear in Trans. Amer. Math. Soc.
- [8] X. He, *The character sheaves on the group compactification*, math.RT/0508068, to appear in Adv. Math.
- [9] X. He, *Minimal length elements in some double cosets of Coxeter groups*, math.RT/0611055.
- [10] X. He and J. F. Thomsen, *Closures of Steinberg fibers in twisted wonderful compactifications*, math.AG/050608, to appear in Trans. Groups.
- [11] X. He and J. F. Thomsen, *Geometry of $B \times B$ -orbit closures in equivariant embeddings*, math.AG/0510088.
- [12] J.-H. Lu and M. Yakimov, *On a class of double cosets in reductive algebraic groups*, International Mathematics Research Notices, **13** (2005), 761-797.
- [13] J.-H. Lu and M. Yakimov, *Group orbits and regular decompositions of Poisson manifolds*, math.SG/0609732.
- [14] G. Lusztig, *Parabolic character sheaves I*, Moscow Math J. **4** (2004), 153-179.
- [15] G. Lusztig, *Parabolic character sheaves II*, Moscow Math J. **4** (2004), 869-896.
- [16] S. Mohrdeiek, *Conjugacy classes of non-connected semisimple algebraic groups*, Dissertation, Univ. Hamburg, 2000.
- [17] P. Slodowy, *Simple Singularities and Simple Algebraic Groups, Lecture Notes in Mathematics, 815*, Springer-Verlag, Berlin, 1980.
- [18] T. A. Springer, *Schubert varieties and generalizations*, In: Representation theories and algebraic geometry (Montreal, PQ, 1997), pp. 413-440, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., 514, Kluwer Acad. Publ., Dordrecht, 1998.
- [19] T. A. Springer, *Intersection cohomology of $B \times B$ -orbit closures in group compactifications*, with an appendix by W. van der Kallen. Special issue in celebration of Claudio Procesi's 60th birthday, J. Algebra **258** (2002), 71-111.
- [20] T. A. Springer, *An extension of Bruhat's Lemma*, preprint, 2006.
- [21] R. Steinberg, *Conjugacy classes in algebraic groups*. Notes by Vinay V. Deodhar, Lect. Notes in Math., Vol. 366. Springer-Verlag, Berlin-New York, 1974.
- [22] R. Steinberg, *Endomorphisms of linear algebraic groups*, Mem. Amer. Math. Soc., **80**, AMS, Providence, RI, 1968.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF HONG KONG, POKFULAM ROAD, HONG KONG
E-mail address: `jhlu@maths.hku.hk`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, SANTA BARBARA, CA 93106, U.S.A.
E-mail address: `yakimov@math.ucsb.edu`